

Optimal Auctions for Asymmetrically Budget Constrained Bidders*

Alexey Malakhov and Rakesh V. Vohra

Department of Finance, Kenan-Flagler School of Business
University of North Carolina
Chapel Hill, NC 27599

Department of Managerial Economics and Decision Sciences
Kellogg School of Management
Northwestern University
Evanston IL 60208, USA

December 2005

Abstract

We consider an environment with a single divisible good and two bidders. The valuations of the bidders are private information but one bidder has a commonly known budget constraint. For this environment we derive the revenue maximizing subsidy free incentive compatible auction. We also examine the case when the budget constraint is private information but bidders must post a bond.

*The research was supported in part by the NSF grant ITR IIS-0121678.

1 Introduction

A seller facing a collection of budget constrained bidders recognizes that such constraints limit the revenue she can generate from the sale of an item. The most obvious limitation is the size of the budgets themselves. They also hinder competition. If one knows that a competitor cannot bid beyond a certain amount, this reduces the bid one needs to place to secure the object. To stimulate competition amongst the bidders, the seller must favor some bidders and handicap others, perhaps by setting reserves or subsidies. While such schemes may increase expected revenues they do so by imposing greater risks on the seller. A reserve, for example, would mean that a trade is sometimes not executed. Finding the right combination of carrots and sticks in this environment to stimulate bidding is therefore an important problem. It is also a difficult one because an analysis must deal with the uncertainty in both a bidders valuations and her budget constraint. The difficulties are not unique to this instance but arise in many environments where the private information of a bidder is multi-dimensional (see, for example, Rochet and Chone (1998) and Rochet and Stole (2003)). For this reason attention has focused on special cases.

Laffont and Robert (1996) assume that valuations are private but that all bidders have the same common knowledge budget constraint. In this environment they derive the subsidy free revenue maximizing Bayesian incentive compatible auction. Since all bidders have the same budget constraint, the analysis sheds no light on the balance between carrots and sticks. The same environment is assumed in Maskin (2000) where he examines constrained efficiency. Again, the symmetry in the budget constraints sheds no light on how the seller should handicap one bidder at the expense of another.

In the same environment as Laffont and Robert, Gaviious, Moldovanu and Sela (2000) compare the revenues to be realized from the standard auction forms. Che and Gale (1996) assume the valuations to be common knowledge but the budgets private information. In this environment they compare the revenues to be realized by the standard auction forms. In addition they examine policies like joint bidding and subsidies and conclude such policies can help to increase the revenue over the standard auction forms. Che and Gale (2000) assume a single buyer with private valuations and a private budget constraint and derive a revenue maximizing, Bayesian

incentive compatible selling mechanism. Since the model involves a single bidder it sheds no light on the issue of competition amongst bidders.

In this paper we consider a problem with two bidders, independent private valuations, bidding for a divisible good. One buyer has no liquidity constraint (the unconstrained bidder), while the other has one. Furthermore, the identity and budget constraint of the constrained bidder is common knowledge. In this environment we derive two mechanisms. One is revenue maximizing, dominant strategy incentive compatible and ex-post individually rational. The second is revenue maximizing, Bayesian incentive compatible and interim individually rational. In addition, we extend the analysis to the case when the budget is private information (but not the identity of the constrained bidder) but the constrained bidder must post a bond. In this case we show how this case reduces to the case when the budget is common knowledge.

All the mechanisms considered are subsidy free. However, as noted by Zheng (2001), offering subsidies may be profitable for the seller. In our model the effect of a subsidy can be computed by increasing the relevant budget constraint and determining if the increase in expected revenue is sufficient to cover the cost of the subsidy.

Two qualitative features of the derived mechanisms differ from the classic optimal mechanism of Myerson (1981) which assumes all bidders are unconstrained. Under a monotone hazard assumption on the distribution of types, Myerson shows that an optimal mechanism will allocate the good to the agent with the highest non-negative virtual value. In the environment considered here, the optimal mechanism may assign a portion of the good to the agent with the lowest virtual value.¹ Furthermore, the good, in some cases, is allocated to an agent with a negative virtual value.

What drives the first difference is the following observation. Suppose the constrained bidder has a value/type i and a budget b . As long as one can offer this bidder a quantity $q \geq b/i$ one can charge her b . Notice that no higher payment from the constrained bidder can be extracted by offering more than b/i . Hence, it may pay to allocate a portion to the unconstrained bidder even though she may have a lower virtual value. For the second

¹Although we assume the good is divisible, one can interpret a fractional allocation as the probability of assigning an indivisible good. In this sense our results carry over to the case of a single indivisible good.

difference, suppose the virtual value of both bidders is negative. It may still be profitable to offer the constrained bidder at least $q \geq b/i$ because one can extract b .

Perhaps the most important lesson to be learnt from the analysis is that the seller must sometimes withhold the good from the unconstrained bidder even if she happens to be the highest bidder. In the event that the unconstrained bidder submits a bid that exceeds the budget of the constrained bidder, the mechanisms derived here, require that the high bid must exceed the budget by a fixed amount in order to secure the good.

The remainder of this paper is divided into three sections. One devoted to the dominant strategy incentive compatible case, the second to the Bayesian incentive compatible case and the third to the case of the budget being private information. In a departure from custom we assume a discrete type space.² This allows us to employ simple arguments involving linear program's to derive the results. It is clear what the continuous analog's of our results are.

2 The Dominant Strategy Incentive Compatible Case

Let $\{1, 2, \dots, m\}$ be the set of possible types, $f_i > 0$ the probability that a bidder is of type i , $F(i) = \sum_{j=1}^i f_j$ and $v_i = [i - \frac{1-F(i)}{f_i}]$ the virtual value of type i . We assume that virtual values are monotone in types (the hazard condition). We use t to denote the type of the unconstrained bidder and s the type of constrained bidder. Let b be the known budget of the constrained bidder.

Our goal is to derive the dominant strategy incentive compatible, ex-post individually rational mechanism that maximizes expected revenue. As is standard, we invoke the revelation principle to restrict attention to direct revelation mechanisms. Let $a(t, s)$ be the allocation to the unconstrained bidder when the profile of reported types is (t, s) and $a_b(t, s)$ the allocation to the constrained bidder when the profile of reported types is (t, s) . The payment that each makes at the profile (t, s) is denoted by $P(t, s)$ and $P_b(t, s)$

²Nevertheless this is consistent with the early work on optimal mechanism design by Harris and Raviv (1981).

respectively.

Dominant strategy incentive compatibility (IC) requires that

$$ta(t, s) - P(t, s) \geq ta(t', s) - P(t', s) \quad \forall t' \neq t,$$

$$sa_b(t, s) - P_b(t, s) \geq sa_b(t, s') - P_b(t, s') \quad \forall s' \neq s.$$

Introducing a dummy type $i = 0$ and invoking the subsidy free assumption allows us to set $P(0, s) = P_b(t, 0) = 0$ and $a(0, s) = a_b(t, 0) = 0$. This enables us to fold the individual rationality (IR) constraint into the (IC) constraints.

The next two results are standard so proofs are omitted.

Theorem 1 *An allocation rule is dominant strategy incentive compatible iff it is monotonic. That is $a(t, s) \geq a(t', s)$ iff. $t \geq t'$ and $a_b(t, s) \geq a_b(t, s')$ iff. $s \geq s'$.*

Theorem 2 *All IC constraints are implied by the following:*

$$ta(t, s) - P(t, s) \geq ta(t - 1, s) - P(t - 1, s) \quad \forall t \leq m,$$

$$ta(t, s) - P(t, s) \geq ta(t + 1, s) - P(t + 1, s) \quad \forall t \leq m - 1$$

$$sa_b(t, s) - P_b(t, s) \geq sa_b(t, s - 1) - P_b(t, s - 1) \quad \forall s \leq m.$$

$$sa_b(t, s) - P_b(t, s) \geq sa_b(t, s + 1) - P_b(t, s + 1) \quad \forall s \leq m - 1$$

The problem of finding the optimal auction is formulated as a linear

program below.

$$\begin{aligned}
[OPT]Z &= \max_{P_b(t,s), P(t,s)} \sum_{t=0}^m \sum_{s=0}^m f_t f_s [P(t,s) + P_b(t,s)] \\
&\text{s.t. } ta(t,s) - P(t,s) \geq ta(t-1,s) - P(t-1,s) \quad \forall t \leq m \\
&ta(t,s) - P(t,s) \geq ta(t+1,s) - P(t+1,s) \quad \forall t \leq m-1 \\
&sa_b(t,s) - P_b(t,s) \geq sa_b(t,s-1) - P_b(t,s-1) \quad \forall s \leq m \\
&sa_b(t,s) - P_b(t,s) \geq sa_b(t,s+1) - P_b(t,s+1) \quad \forall s \leq m-1 \\
&a(t,s) \geq a(t-1,s) \quad \forall t \leq m, s \\
&a_b(t,s) \geq a_b(t,s-1) \quad \forall s \leq m, t \\
&P_b(t,s) \leq b \quad \forall t, s \\
&a(t,s) + a_b(t,s) \leq 1 \quad \forall t, s \\
&a(t,s), a_b(t,s) \geq 0 \quad \forall t, s
\end{aligned}$$

Fixing the a 's, a_b 's and the type t of the unconstrained bidder, $P_b(t, \cdot)$ can be expressed as a shortest path length in an appropriate network.

In this network there is one vertex for each type s of the constrained bidder, including the dummy type. The vertex corresponding to the dummy type is called the source vertex. For each type/vertex $i \geq 1$ there is a directed edge $(0, i)$ of length b (only the edge from source to vertex i is depicted). Between the source and vertex $\{1\}$ there are two parallel edges. One of length $a_b(t, 1)$ and the other of length b .

Between i and $i+1$ there are two edges. One directed from i to $i+1$ of length $(i+1)[a_b(t, i+1) - a_b(t, i)]$. Monotonicity of the allocation rule implies this edge has non-negative length. The other edge is directed from $i+1$ to i and of length $i[a_b(t, i) - a_b(t, i+1)] \leq 0$. A portion of the network is depicted in Figure 1 (the edge of length b from the source to vertex 1 is not depicted.)




Figure 1

The edges directed from left to right correspond to ‘downward’ IC constraints while the edges in the opposite directions correspond to ‘upward’ IC constraints. An implication of monotonicity of the a_b ’s is that there are no negative cycles in the network.

The largest value that $P_b(t, i)$ can take is the length of the shortest path from the source to vertex i . The absence of negative length cycles means these shortest path lengths are well defined. In this network there are at most two candidates for the shortest path from the source to $i < m$. One from left to right and one from right to left. Let $L_i^b(t)$ be the length of the path from node 0 to node i that is from left to right, and $R_i^b(t)$ be the length of the path from node 0 to node i that is from right to left. In particular $L_0^b(t) = 0$, $L_i^b(t) = \sum_{j=0}^{i-1} (j+1)[a_b(t, j+1) - a_b(t, j)] \forall i > 0$. Similarly $R_i^b(t) = b - \sum_{j=i}^{m-1} j[a_b(t, j+1) - a_b(t, j)] \forall i < m$ and $R_m^b(t) = b$. Clearly, $P_b(t, i) = \min(L_i^b(t), R_i^b(t))$.

A similar network can be constructed to determine $P(i, s)$. It will be identical to the one in Figure 1 except the edges of length b from the source to each vertex i will be absent. It is easy to see from an identical analysis

that

$$P(i, s) = \sum_{j=0}^{i-1} (j+1)[a(j+1, s) - a(j, s)].$$

We can rewrite [OPT] as

$$\begin{aligned} Z &= \max_{P_b(t,s), P(t,s)} \sum_{t=0}^m \sum_{s=0}^m f_t f_s [P(t, s) + P_b(t, s)] \\ \text{s.t. } P_b(t, s) &= \min\{L_s^b(t), R_s^b(t)\} \quad \forall t, s \\ P(t, s) &= \sum_{j=0}^{t-1} (j+1)[a(j+1, s) - a(j, s)] \quad \forall t, s \\ a(t, s) &\geq a(t-1, s) \quad \forall t \leq m, s \\ a_b(t, s) &\geq a_b(t, s-1) \quad \forall s \leq m, t \\ a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\ a(t, s), a_b(t, s) &\geq 0 \quad \forall t, s \end{aligned}$$

Straightforward algebra shows that

$$\sum_{t=0}^m \sum_{s=0}^m f_t f_s P(t, s) = \sum_{t=0}^m \sum_{s=0}^m f_t f_s \left(\sum_{j=0}^{t-1} (j+1)[a(j+1, s) - a(j, s)] \right) = \sum_{s=0}^m f_s \left[\sum_{t=0}^m f_t v_t a(t, s) \right].$$

Substituting this into [OPT] to:

$$\begin{aligned} Z &= \max_{P_b(t,s), a(t,s)} \sum_{s=0}^m f_s \left[\sum_{t=0}^m f_t v_t a(t, s) \right] + \sum_{t=0}^m \sum_{s=0}^m f_t f_s P_b(t, s) \\ \text{s.t. } P_b(t, s) &= \min\{L_s^b(t), R_s^b(t)\} \quad \forall t, s \\ a(t, s) &\geq a(t-1, s) \quad \forall t \leq m, s \\ a_b(t, s) &\geq a_b(t, s-1) \quad \forall s \leq m, t \\ a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\ a(t, s), a_b(t, s) &\geq 0 \quad \forall t, s \end{aligned}$$

Monotonicity of the a_b 's implies that $L_i^b(t) \leq L_{i+1}^b(t)$ and $b \geq R_i^b(t) \geq R_{i+1}^b(t)$ for all t . Hence the budget constraint is already incorporated into the formulation.

Consider now the following relaxation of $[OPT]$ which we call $[rOPT]$.

$$\begin{aligned}
& \max_{P_b(t,s), a(t,s)} \sum_{s=0}^m f_s \left[\sum_{t=0}^m f_t v_t a(t,s) \right] + \sum_{t=0}^m \sum_{s=0}^m f_t f_s P_b(t,s) \\
& \text{s.t. } P_b(t,s) \leq L_s^b(t) \quad \forall t, s \\
& P_b(t,s) \leq b \quad \forall t, s \\
& a(t,s) \geq a(t-1,s) \quad \forall t \leq m, s \\
& a_b(t,s) \geq a_b(t,s-1) \quad \forall s \leq m, t \\
& a(t,s) + a_b(t,s) \leq 1 \quad \forall t, s \\
& a(t,s), a_b(t,s) \geq 0 \quad \forall t, s
\end{aligned}$$

Lemma 1 *There is an optimal solution to $[rOPT]$ such that $L_s^b(t) \leq b$ for all s and t .*

Proof

Amongst all optimal solutions to $[rOPT]$ (this forms a compact set) pick one that minimizes $D = \sum_{t=1}^m \max\{L_m^b(t) - b, 0\}$. If $D = 0$, monotonicity of the a_b 's implies that $L_s^b(t) \leq L_m^b(t) \leq b$ for all s and t and the proof is complete. Suppose therefore that $D > 0$. Hence there is a t such that $L_m^b(t) > b$. Let k be the smallest index such that $a_b(t,k) = a_b(t,k+1) = \dots = a_b(t,m)$. Notice that $a_b(t,m) > 0$. Construct a new solution by reducing $a_b(s,t)$ for all $s \geq k$ by $\epsilon > 0$. Observe that this does not influence $P_b(t,s)$ for all s . For $s < k$ we have that $a_b(t,s) < a_b(t,k)$ and so for ϵ sufficiently small the new solution is still monotonic in the allocation variables. Thus, we have a new solution that is feasible, optimal but with a value of D that is reduced, a contradiction. ■

In view of lemma 1 and the monotonicity of the a 's we can write $[rOPT]$

as

$$\begin{aligned}
& \max_{\{a(t,s), a_b(t,s)\}} \sum_t \sum_s f_t f_s [v_t a(t,s) + v_s a_b(t,s)] & (P) \\
& \text{s.t. } a(t,s) \leq a(t+1,s) \quad \forall t, s \\
& \quad a_b(t,s) \leq a_b(t,s+1) \quad \forall t, s \\
& \quad a(t,s) + a_b(t,s) \leq 1 \quad \forall t, s \\
& \quad a(t,s), a_b(t,s) \geq 0 \quad \forall t, s \\
& \sum_{s=0}^{m-1} (s+1) [a_b(t,s+1) - a_b(t,s)] \leq b \quad \forall t
\end{aligned}$$

Notice that any solution to this last program is feasible for $[OPT]$ as well. This is because $L_s^b(t) \leq b \Rightarrow L_s^b(t) \leq R_s^b(t)$.

2.1 A Relaxation of problem (P)

We relax problem (P) by removing the monotonicity constraints on a .

$$\begin{aligned}
& \max_{\{a(t,s), a_b(t,s)\}} \sum_t \sum_s f_t f_s [v_t a(t,s) + v_s a_b(t,s)] & (P_1) \\
& \text{s.t. } a_b(t,s) \leq a_b(t,s+1) \quad \forall t, s \\
& \quad a(t,s) + a_b(t,s) \leq 1 \quad \forall t, s \\
& \quad a(t,s), a_b(t,s) \geq 0 \quad \forall t, s \\
& \sum_{s=0}^{m-1} (s+1) [a_b(t,s+1) - a_b(t,s)] \leq b \quad \forall t
\end{aligned}$$

We will solve (P_1) and it will be verified that the solution satisfies the omitted monotonicity constraints. Thus the relaxation is exact.

Problem (P_1) itself can be decomposed into a collection of independent sub-problems indexed by the type t of the unconstrained bidder:

$$\begin{aligned}
& \max_{\{a(t,s), a_b(t,s)\}} \sum_s f_s [v_t a(t,s) + v_s a_b(t,s)] & (P_t) \\
& \text{s.t. } a_b(t,s) \leq a_b(t,s+1) \quad \forall s \\
& \quad a(t,s) + a_b(t,s) \leq 1 \quad \forall s \\
& \quad m a_b(t,m) - \sum_{j=1}^{m-1} a_b(t,j) \leq b \quad \forall t
\end{aligned}$$

We divide the solution of (P_t) into two cases.

2.2 Solution of (P_t) when $v_t \geq 0$

We can deal with the monotonicity constraints on the a_b 's by a change of variables. Let $a_b(t, s) = \sum_{i=1}^s \Delta(t, i)$ where $\Delta(t, i) \geq 0$ for all i . For convenience set $w_s = \sum_{j=s}^m f_j v_j$ for all s . Replacing the a_b 's by Δ 's in problem (P_t) yields:

$$\begin{aligned} \max_{\{a(t,s), \Delta(s,t)\}} & \sum_{s=1}^m f_s v_t a(t, s) + \sum_{s=1}^m w_s \Delta(t, s) & (LP_t) \\ \text{s.t.} & a(t, s) + \sum_{i=1}^s \Delta(t, i) \leq 1 \quad \forall s \\ & \sum_{s=1}^m s \Delta(t, s) \leq b \\ & a(t, s), \Delta(t, s) \geq 0 \quad \forall s \end{aligned}$$

It is easy to see that there is an optimal solution to (LP_t) such that $a(t, s) + \sum_{i=1}^s \Delta(t, i) = 1$ for all s . If not, i.e. $a(t, s) + \sum_{i=1}^s \Delta(t, i) < 1$ for some s , increase $a(t, s)$. Since $v_t \geq 0$ the objective function cannot decrease. Given this we can use the equation $a(t, s) + \sum_{i=1}^s \Delta(t, i) = 1$ to substitute out the $a(t, s)$ variables. Specifically $a(t, s) = 1 - \sum_{i=1}^s \Delta(t, i)$ for all s . Therefore (LP_t) becomes:

$$\begin{aligned} \max_{\{a(t,s), \Delta(s,t)\}} & \sum_{s=1}^m f_s v_t [1 - \sum_{i=1}^s \Delta(t, i)] + \sum_{s=1}^m w_s \Delta(t, s) & (LP_t) \\ \text{s.t.} & \sum_{i=1}^s \Delta(t, i) \leq 1 \quad \forall s \\ & \sum_{s=1}^m s \Delta(t, s) \leq b \\ & \Delta(t, s) \geq 0 \quad \forall s \end{aligned}$$

A second observation is the constraints $\sum_{i=1}^s \Delta(t, i) \leq 1$ for all $s = 1, \dots, m-1$ are all implied by the constraint $\sum_{i=1}^m \Delta(t, i) \leq 1$. This allows us to reduce

(LP_t) to

$$\begin{aligned}
\max_{\{a(t,s), \Delta(s,t)\}} & \sum_{s=1}^m f_s v_t [1 - \sum_{i=1}^s \Delta(t, i)] + \sum_{s=1}^m w_s \Delta(t, s) & (\text{LP}_t) \\
\text{s.t.} & \sum_{i=1}^m \Delta(t, i) \leq 1 \\
& \sum_{s=1}^m s \Delta(t, s) \leq b \\
& \Delta(t, s) \geq 0 \quad \forall s
\end{aligned}$$

Simplifying the objective function and introducing slack variables u_1 and u_2 to make all constraints hold at equality produces:

$$\begin{aligned}
\max_{\{\Delta(t,s)\}} & \sum_{s=1}^m [\sum_{i=s}^m f_i (v_i - v_t)] \Delta(t, s) + \sum_{s=1}^m f_s v_t \\
\text{s.t.} & \sum_{s=1}^m \Delta(t, s) + u_1 = 1 \\
& \sum_{s=1}^m s \Delta(t, s) + u_2 = b \\
& \Delta(t, s), u_i \geq 0 \quad \forall s, i
\end{aligned}$$

The extreme points of this last linear program fall into one of three categories because it has only two constraints.

1. Exactly one i such that $\Delta(t, i) > 0$, $u_1 > 0$, $u_2 = 0$.
2. Exactly one i such that $\Delta(t, i) > 0$, $u_1 = 0$, $u_2 > 0$.
3. Exactly one pair (p, q) with $p \neq q$, $p \leq b \leq q$, $\Delta(t, p), \Delta(t, q) > 0$ and $u_1 = u_2 = 0$.

From each category we choose the extreme point with largest objective function value. The optimal solution must be the extreme point from the three chosen with largest objective function value. The best extreme point of each type is listed below.

1. **Category 1:**

Let $r^1 = \arg \max_{s \geq b} \frac{b \sum_{i=s}^m f_i(v_i - v_t)}{s}$.

Set $\Delta(t, r^1) = \frac{b}{r_1}$, $u_1 = 1 - \frac{b}{r_1}$ and $u_2 = 0$.

The objective function value of this solution is $b \frac{\sum_{i=r^1}^m f_i(v_i - v_t)}{s} + \sum_{s=1}^m f_s v_t$.

2. **Category 2:**

Let $r^2 = \arg \max_{s \leq b} [\sum_{i=s}^m f_i(v_i - v_t)]$. In fact, by the monotone hazard condition, when $b \leq t$, $r_2 = b$ and when $b > t$, $r_2 = t$.

Set $\Delta(t, r^2) = 1$, $u_1 = 0$ and $u_2 = b - r^2$.

The objective function value of this solution is $\sum_{i=r^2}^m f_i(v_i - v_t) + \sum_{s=1}^m f_s v_t$.

3. **Category 3:** Let

$$(p^3, q^3) = \arg \max_{p \leq b \leq q} \left[\frac{(q-b)}{(q-p)} \sum_{i=p}^m f_i(v_i - v_t) + \frac{(b-p)}{(q-p)} \sum_{i=q}^m f_i(v_i - v_t) \right].$$

Set $\Delta(t, p^3) = \frac{(q^3-b)}{(q^3-p^3)}$, $\Delta(t, q^3) = \frac{(b-p^3)}{(q^3-p^3)}$ and $u_1 = u_2 = 0$.

In fact we can pin down the category 3 solution even further. Suppose first that $b \leq t$. Then, by the monotone hazard condition, $\sum_{i=q}^m f_i(v_i - v_t)$ is maximized when $q = t$ and $\sum_{i=p}^m f_i(v_i - v_t)$ is maximized when $p = b$. In this case the category 3 solution is $\Delta(t, p^3) = 1$ and $\Delta(t, q^3) = 0$, i.e. a category 2 solution. A similar argument applies in the case when $b > t$.

Therefore, only category 1 and 2 solutions apply.

Theorem 3 *If $b > t$ then $\Delta(t, t) = 1$ is optimal. If $b \leq t$ then $\Delta(t, r^1) = \frac{b}{r_1}$ is optimal.*

Proof

Suppose first $b > t$. By the monotone hazard condition, $\frac{\sum_{i=s}^m f_i(v_i - v_t)}{s}$ is non-increasing for $s \geq t$. Hence

$$\arg \max_{s \geq b} \frac{b \sum_{i=s}^m f_i(v_i - v_t)}{s} = b.$$

In this case the objective function value of the category 1 solution is $\sum_{s=b}^m f_s(v_s - v_t)$. However the category 2 solution has objective function value $\sum_{s=t}^m f_s(v_s - v_t)$, which is larger.

Now suppose $b \leq t$. The category 2 solution has objective function value $\sum_{s=b}^m f_s(v_s - v_t)$. But this is bounded above by $\max_{s \geq b} \frac{b \sum_{i=s}^m f_i(v_i - v_t)}{s}$, i.e., the category 1 solution is optimal. ■

To summarize:

1. If $b > t$ then $a_b(t, s) = 1, a(t, s) = 0$ for all $s \geq t$ and $a_b(t, s) = 0, a(t, s) = 1$ otherwise.
2. If $b \leq t$ then $a_b(t, s) = \frac{b}{r^1}, a(t, s) = 1 - \frac{b}{r^1}$ for all $s \geq r^1$ and $a_b(t, s) = 0, a(t, s) = 1$ otherwise.

Using the expressions for for the payment variables in terms of path lengths we can compute the payments bidders must make. The allocations are summarized in the table below.

$v_t \geq 0$	$s \geq r^1$	$b \leq s \leq r^1$	$s \leq b - 1$
$t \geq b$	$a(t, s) = (1 - \frac{b}{r^1})$ $a_b(t, s) = \frac{b}{r^1}$	$a(t, s) = 1$ $a_b(t, s) = 0,$	$a(t, s) = 1$ $a_b(t, s) = 0$
$t \leq b - 1$	$a(t, s) = 0$ $a_b(t, s) = 1$	$a(t, s) = 0,$ $a_b(t, s) = 1$	usual auction rules

Examination of the table shows that a satisfies the omitted monotonicity constraint.

2.3 Solution of (P_t) when $v_t < 0$

In problem (P_t) if $v_t < 0$ then $a(t, i) = 0$ for all i in any optimal solution. Note that if $v_i < 0$ for some i we cannot conclude that $a_b(t, i) = 0$ in every optimal solution because of the budget constraint. Problem (P_t) becomes

$$\begin{aligned}
& \max_{\{\Delta(s,t)\}} \sum_{i=1}^m w_s^t \Delta(t, s) && (LP'_t) \\
& \text{s.t.} \quad \sum_{i=1}^s \Delta(t, i) \leq 1 \quad \forall s \\
& \quad \quad \sum_{s=1}^m s \Delta(t, s) \leq b \\
& \quad \quad \Delta(t, s) \geq 0 \quad \forall s
\end{aligned}$$

The similarity of (LP'_t) to (LP_t) permits an identical analysis whose details are omitted. The conclusions are summarized below. Let $h^1 = \arg \max_{s \geq b} \frac{b \sum_{i=s}^m w_i}{s}$.

1. If $w_1 \geq b \frac{w_{h^1}}{h^1}$ then $a_b(t, s) = 1, a(t, s) = 0$ for all $s \geq 1$ and $a_b(t, s) = 0, a(t, s) = 0$ otherwise.
2. If $w_1 < b \frac{w_{h^1}}{h^1}$ then $a_b(t, s) = \frac{b}{h^1}, a(t, s) = 0$ for all $s \geq h^1$ and $a_b(t, s) = 0, a(t, s) = 0$ otherwise.

Notice that the a satisfy the omitted monotonicity constraint.

3 The Bayesian Incentive Compatible Case

Here we derive the revenue maximizing auction that is Bayesian incentive compatible and interim individually rational. Let \mathcal{A}_i^b be the **expected** quantity of the good that the constrained agent receives when she reports type i and all other agents report truthfully. Let p_i^b be the **expected** payment of the constrained agent who reports type i . Similarly, \mathcal{A}_i and p_i are the expected allocations and payments for the unconstrained agent who reports type i . Notice that $\mathcal{A}_i^b = \sum_t f_t a_b(t, i)$ and $p_i^b = \sum_{t=1}^m f_t P_b(t, i)$ for all i . Similar expressions hold for \mathcal{A}_i and p_i^u . Bayesian incentive compatibility (BIC) for the constrained agent requires:

$$i\mathcal{A}_i^b - p_i^b \geq i\mathcal{A}_j^b - p_j^b \quad \forall j \neq i.$$

A similar inequality holds for the unconstrained agent.

Introducing a dummy type $i = 0$ with $p_0^u, p_0^b = 0$ and $\mathcal{A}_0^b, \mathcal{A}_0^u = 0$ allows us to fold the interim individual rationality constraint into the (BIC) constraints.

From now on statements about \mathcal{A}_i and p_i are to be read as applying to \mathcal{A}_i^b and p_i^b as well.

The next two results are standard.

Theorem 4 *An allocation rule that is incentive compatible iff it is monotonic. That is $r \leq s$ iff $\mathcal{A}_r \leq \mathcal{A}_s$ for all $k = 1, \dots, m$.*

Theorem 5 *All (BIC) constraints are implied by the following:*

$$i\mathcal{A}_i - p_i \geq i\mathcal{A}_{i-1} - p_{i-1} \quad \forall i = 1, \dots, m$$

$$i\mathcal{A}_i - p_i \geq i\mathcal{A}_{i+1} - p_{i+1} \quad \forall i = 1, \dots, m - 1$$

We can formulate the problem of finding the optimal auction as:

$$\begin{aligned}
[BOPT] \quad Z &= \max_{p_i, p_i^b} \sum_{s,t}^m f_t f_s (p_t + p_s^b) \\
\text{s.t.} \quad i\mathcal{A}_i - p_i &\geq i\mathcal{A}_{i-1} - p_{i-1} \quad \forall i = 1, \dots, m \\
i\mathcal{A}_i - p_i &\geq i\mathcal{A}_{i+1} - p_{i+1} \quad \forall i = 1, \dots, m-1 \\
\mathcal{A}_i &= \sum_s f_s a(i, s) \quad \forall i \\
\mathcal{A}_i^b &= \sum_t f_t a_b(t, i) \quad \forall i \\
p_i &= \sum_{s=1}^m f_t P(i, s) \quad \forall i \\
p_i^b &= \sum_{t=1}^m f_t P_b(t, i) \quad \forall i \\
a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\
p_i^b &\leq b \quad \forall i
\end{aligned}$$

The last constraint requires that the *expected* payment not exceed b .

As in the previous case there is a representation of the BIC constraints that allows one to interpret the P_i 's as shortest path lengths in a network. Specifically,

$$p_s^b = \min\left\{\sum_{j=0}^{s-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b), b - \sum_{j=i}^{m-1} j[\mathcal{A}_{j+1}^b - \mathcal{A}_j^b]\right\}$$

and

$$p_t = \sum_{j=0}^{t-1} (j+1)(\mathcal{A}_{j+1} - \mathcal{A}_j)$$

for all s, t . An analysis identical to the dominant strategy case allows us to write [BOPT] as

$$\begin{aligned}
\max \sum_{s,t} & [f_t(\sum_{j=0}^{t-1} (j+1)(\mathcal{A}_{j+1} - \mathcal{A}_j)) + f_s(\sum_{j=0}^{s-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b))] \\
\text{s.t.} \quad & \sum_{j=0}^{i-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b) \leq b \quad \forall i > 0 \\
& \mathcal{A}_1 \leq \dots \leq \mathcal{A}_m \\
& \mathcal{A}_i = \sum_s f_s a(i, s) \quad \forall i \\
& \mathcal{A}_i^b = \sum_t f_t a_b(t, i) \quad \forall i \\
& a(t, s) + a_b(t, s) \leq 1 \quad \forall t, s \\
& a(t, s), a_b(t, s) \geq 0 \quad \forall t, s
\end{aligned}$$

Collecting like terms in the objective function together shows that

$$\sum_{s,t} [f_t(\sum_{j=0}^{t-1} (j+1)(\mathcal{A}_{j+1} - \mathcal{A}_j)) + f_s(\sum_{j=0}^{s-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b))] = \sum_{s,t} [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s].$$

The constraints $\sum_{j=0}^{i-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b) \leq b$ for all $i > 0$ can be rewritten to read $i\mathcal{A}_i^b \leq b + \sum_{j=1}^{i-1} \mathcal{A}_j^b$. For each pair i label the constraint $i\mathcal{A}_i^b \leq b + \sum_{j=1}^{i-1} \mathcal{A}_j^b$ as C_i .

Problem [BOPT] becomes:

$$\begin{aligned}
Z = \max \sum_{s,t} & [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s] \\
\text{s.t.} \quad & i\mathcal{A}_i^b \leq b + \sum_{j=1}^{i-1} \mathcal{A}_j^b \quad \forall i \\
& 0 \leq \mathcal{A}_1 \leq \dots \leq \mathcal{A}_m \\
& \mathcal{A}_i = \sum_s f_s a(i, s) \quad \forall i \\
& \mathcal{A}_i^b = \sum_t f_t a_b(t, i) \quad \forall i \\
& a(t, s) + a_b(t, s) \leq 1 \quad \forall t, s \\
& a(t, s), a_b(t, s) \geq 0 \quad \forall t, s
\end{aligned}$$

Lemma 2 *Suppose in some optimal solution to [BOPT] one or more of the C_i are binding. Let s^* be the smallest index for which C_i is binding. Then there is an optimal solution to [BOPT] such that*

$$\mathcal{A}_m^b = \dots = \mathcal{A}_{s^*}^b.$$

Proof

Since C_{s^*} is binding we have $s^* \mathcal{A}_{s^*}^b = b + \sum_{j=1}^{s^*-1} \mathcal{A}_j^b$. From C_{s^*+1} we have

$$(s^* + 1) \mathcal{A}_{s^*+1}^b \leq b + \sum_{j=1}^{s^*} \mathcal{A}_j^b = \mathcal{A}_{s^*}^b + s^* \mathcal{A}_{s^*}^b = (s^* + 1) \mathcal{A}_{s^*}^b.$$

But $\mathcal{A}_{s^*+1}^b \geq \mathcal{A}_{s^*}^b$, which implies that $\mathcal{A}_{s^*+1}^b = \mathcal{A}_{s^*}^b$. Thus C_{s^*+1} is binding and we repeat the argument for index $s^* + 2$ and so on. \blacksquare

Suppose we knew the critical index s^* from the lemma. Consider the following optimization problem [BOPT(s^*)]:

$$\begin{aligned} Z(s^*) &= \max \sum_{s,t} [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s] \\ \text{s.t. } \mathcal{A}_1^b &\leq \dots \leq \mathcal{A}_{s^*}^b = \dots = \mathcal{A}_m^b \\ \mathcal{A}_1 &\leq \dots \leq \mathcal{A}_m \\ \mathcal{A}_i &= \sum_s f_s a(i, s) \quad \forall i \\ \mathcal{A}_i^b &= \sum_t f_t a_b(t, i) \quad \forall i \\ a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\ a(t, s), a_b(t, s) &\geq 0 \quad \forall s, t \end{aligned}$$

It is clear that $Z = Z(s^*)$. We will show that there is an optimal solution to [BOPT(s^*)] such that $a_b(t, s) = a_b(t, s^*)$ for all t and $s \geq s^*$. We do this

by examining a relaxation of problem $[BOPT(s^*)]$ called $[ROPT(s^*)]$.

$$\begin{aligned}
Z^r(s^*) &= \max_{s,t} \sum [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s] \\
&\text{s.t. } \mathcal{A}_{s^*}^b = \dots = \mathcal{A}_m^b \\
\\
\mathcal{A}_i &= \sum_s f_s a(i, s) \quad \forall i \\
\mathcal{A}_i^b &= \sum_t f_t a_b(t, i) \quad \forall i \\
a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\
a(t, s), a_b(t, s) &\geq 0 \quad \forall s, t
\end{aligned}$$

The relaxation is obtained by removing the monotonicity constraints on \mathcal{A}_j for all j , the monotonicity constraints on \mathcal{A}_j^b for $j = 1, \dots, s^* - 1$. We show that there is an optimal solution to the relaxation that satisfies the omitted constraints, so making the relaxation exact.

3.1 Solving $[ROPT(s^*)]$

The Lagrangean dual of $[ROPT(s^*)]$ is problem (P_λ) , shown below.

$$\begin{aligned}
Z_\lambda(s^*) &= \max_{s,t} \sum [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s] + \sum_{j=s^*}^{m-1} \lambda_j (\mathcal{A}_{j+1}^b - \mathcal{A}_j^b) \\
&\text{s.t. } \mathcal{A}_i = \sum_s f_s a(i, s) \quad \forall i \\
\mathcal{A}_i^b &= \sum_t f_t a_b(t, i) \quad \forall i \\
a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\
a(t, s), a_b(t, s) &\geq 0 \quad \forall s, t
\end{aligned}$$

By the duality theorem, $Z^r(s^*) = \min_\lambda Z_\lambda(s^*)$.

The objective function of (P_λ) can be written as

$$\sum_{t=1}^m f_t \mathcal{A}_t v_t + \sum_{s=1}^{s^*-1} f_s \mathcal{A}_s^b v_s + \sum_{s=s^*}^m f_s \left[v_s + \frac{\lambda_{s-1}}{f_s} - \frac{\lambda_s}{f_s} \right] \mathcal{A}_s^b$$

where $\lambda_{s^*-1} = \lambda_m = 0$. Substituting $\mathcal{A}_i = \sum_s f_s a(i, s)$ and $\mathcal{A}_i^b = \sum_t f_t a_b(t, i)$ into this expression yields:

$$\sum_{t=1}^m \sum_{s=1}^m f_t v_t f_s a(t, s) + \sum_{s=1}^{s^*-1} \sum_{t=1}^m f_s v_s f_t a_b(t, s) + \sum_{s=s^*}^m \sum_{t=1}^m f_s \left[v_s + \frac{\lambda_{s-1}}{f_s} - \frac{\lambda_s}{f_s} \right] f_t a_b(t, s).$$

To simplify, let $h_s(\lambda) = v_s + \frac{\lambda_{s-1}}{f_s} - \frac{\lambda_s}{f_s}$ for $s \geq s^*$. Therefore

$$Z_\lambda(s^*) = \max \sum_{t=1}^m \sum_{s=1}^m f_t v_t f_s a(t, s) + \sum_{s=1}^{s^*-1} \sum_{t=1}^m f_s v_s f_t a_b(t, s) + \sum_{s=s^*}^m \sum_{t=1}^m f_s h_s(\lambda) f_t a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1 \quad \forall t, s$
 $a(t, s), a_b(t, s) \geq 0 \quad \forall s, t$

This decomposes into a collection of subproblems, one subproblem for each profile (t, s) of types. When $s \leq s^* - 1$ the subproblem is

$$g_\lambda(t, s) = \max v_t a(t, s) + v_s a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1$
 $a(t, s), a_b(t, s) \geq 0$

and when $s \geq s^*$ it is

$$g_\lambda(t, s) = \max v_t a(t, s) + h_s(\lambda) a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1$
 $a(t, s), a_b(t, s) \geq 0$

It is easy to see that for $s \leq s^* - 1$ that $g_\lambda(t, s) = \max\{v_t, v_s\}$ and when $s \geq s^*$, $g_\lambda(t, s) = \max\{v_t, h_s(\lambda)\}$.

Therefore $Z_\lambda(s^*) = \sum_{t,s} f_t f_s g_\lambda(t, s)$. This allows us to formulate $\min_\lambda Z_\lambda(s^*)$ as a linear program (LP_λ):

$$\min \sum_{t,s} f_t f_s W(t, s)$$

s.t. $W(t, s) \geq v_t \quad \forall t$
 $W(t, s) \geq v_s \quad \forall s \leq s^* - 1$
 $W(t, s) \geq h_s(\lambda) \quad \forall s \geq s^*$
 $\lambda_{s^*-1}, \lambda_m = 0$

Lemma 3 *There is an optimal solution (W^*, λ^*) of (LP_λ) such that $v_{s^*-1} \leq h_{s^*}^* \leq \dots \leq h_m^*$ where $h_s^* = v_s + \frac{\lambda_{s-1}^*}{f_s} - \frac{\lambda_s^*}{f_s}$ for $s \geq s^*$.*

Proof

Denote the discrepancy of an optimal solution λ^* to (LP_λ) by $\max\{v_{s^*-1} - h_{s^*}^*, 0\} + \sum_{i \geq s^*+1} \max\{h_{i-1}^* - h_i^*, 0\}$. Amongst all optimal solutions to (LP_λ) pick the one that has the smallest discrepancy. If the discrepancy is zero, we are done. So, suppose not.

Case 1: There exists at least one $j \geq s^* + 1$, such that $h_{j-1}^* > h_j^*$. If there exist more than one j such that $h_{j-1}^* > h_j^*$, choose the largest j , for which $h_{j-1}^* > h_j^*$.

Let l be the largest index such that $h_j^* = h_{j+1}^* = \dots = h_l^* < h^*l + 1$.³ Suppose $l < m$.

We construct a contradiction by considering a new set of $\{\lambda'_i\}_{i=0}^m$, such that

$$\begin{aligned} \lambda'_{j-1} &= \lambda_{j-1}^* + \varepsilon, \\ \lambda'_i &= \lambda_i^* - \varepsilon \quad \forall i \in [j, l], \\ \lambda'_i &= \lambda_i^* \quad \forall i \in [1, j-2] \cup [l+1, m]. \end{aligned}$$

Denote $h'_i = i - \frac{1-F(i)}{f_i} - \frac{\lambda'_i}{f_i} + \frac{\lambda'_{i-1}}{f_i}$ for all i . This change results in the following changes to the values of $\{h_i^*\}_{i=0}^m$:

$$\begin{aligned} h'_{j-1} &= h_{j-1}^* - \frac{\varepsilon}{f_{j-1}}, \\ h'_j &= h_j^* + 2\frac{\varepsilon}{f_j}, \\ h'_{l+1} &= h_{l+1}^* - \frac{\varepsilon}{f_{l+1}}, \\ h'_i &= h_i^* \quad \forall i \notin \{j-1, j, l+1\}. \end{aligned}$$

Denote the change in the (LP_λ) problem objective function from changing λ^* to λ' by $\Delta Z_\lambda(s^*)$. Consider the pairs (t, s) , for which $W(t, s)$ are affected by this change.

For $\varepsilon > 0$ sufficiently small, decreasing h_{j-1}^* by $\frac{\varepsilon}{f_{j-1}}$ affects $W(s, t)$ only if $v_t < h_{j-1}^*$. Similarly the decrease of h_{l+1}^* by $\frac{\varepsilon}{f_{l+1}}$ affects $W(t, s)$ if $v_t < h_{l+1}^*$.

³It is possible that $l = j$.

For $\varepsilon > 0$ sufficiently small, increasing h_j^* by $2\frac{\varepsilon}{f_j}$ affects $W(t, s)$ only if $v_t \leq h_j$.

Therefore the change in the objective function $\Delta Z_\lambda(s^*)$ is:

$$\Delta Z_\lambda(s^*) \leq 2\varepsilon[\sum\{f_t : v_t \leq h_j^*\}] - \varepsilon[\sum\{f_t : v_t < h_{j-1}^*\}] - \varepsilon\sum\{f_t : v_t < (h_{l+1}^*)\}.$$

Hence $\Delta Z_\lambda(s^*) \leq 0$, and we conclude that λ' is also an optimal solution to (LP_λ) .

Computing the change in discrepancy from λ^* to λ' we observe that $\max\{h_{j-2} - h_{j-1}, 0\}$ can increase by at most ε/f_{j-1} , the term $\max\{h_{j-1} - h_j, 0\}$ goes down by $\varepsilon/f_{j-1} + 2\varepsilon/f_j$, the term $\max\{h_j - h_{j+1}, 0\}$ goes up by $2\varepsilon/f_j$, and the term $\max\{h_l - h_{l+1}, 0\}$ goes down by ε/f_{l+1} . The contribution to discrepancy from other terms is unchanged.⁴ Notice that the discrepancy changes by $\varepsilon/f_{j-1} - (\varepsilon/f_{j-1} + 2\varepsilon/f_j) + 2\varepsilon/f_j - \varepsilon/f_{l+1} < 0$, contradicting our choice of λ^* as the one with the smallest discrepancy.

Now suppose that $l = m$. This implies that $h_j^* = h_{j+1}^* = \dots = h_m^*$.⁵

We construct a contradiction by considering a new set of $\{\lambda'_i\}_{i=0}^m$, such that

$$\begin{aligned}\lambda'_i &= \lambda_i^* + \varepsilon \quad \forall i \in [j-1, m-1], \\ \lambda'_i &= \lambda_i^* \quad \forall i \notin [j-1, m-1].\end{aligned}$$

Denote $h'_i = i - \frac{1-F(i)}{f_i} - \frac{\lambda'_i}{f_i} + \frac{\lambda'_{i-1}}{f_i}$ for all i . This change results in the following changes to the values of $\{h_i^*\}_{i=0}^m$:

$$\begin{aligned}h'_{j-1} &= h_{j-1}^* - \frac{\varepsilon}{f_{j-1}}, \\ h'_m &= h_m^* + \frac{\varepsilon}{f_m}, \\ h'_i &= h_i^* \quad \forall i \notin \{j-1, m\}.\end{aligned}$$

For $\varepsilon > 0$ sufficiently small, decreasing h_{j-1}^* by $\frac{\varepsilon}{f_{j-1}}$ affects $W(t, s)$ only if $v_t < h_{j-1}^*$. Increasing h_m^* by $\frac{\varepsilon}{f_m}$ affects $W(t, s)$ only if $v_t \leq h_m^*$.

The change in the objective function $\Delta Z_\lambda(s^*)$ is:

$$\Delta Z_\lambda(s^*) \leq \varepsilon[\sum\{f_t : v_t \leq h_m^*\}] - \varepsilon\sum\{f_t : v_t < h_{j-1}^*\} \leq 0.$$

⁴the term $\max\{h_j - h_{j+1}, 0\}$ is also unchanged for a small enough $\varepsilon > 0$.

⁵It is possible that $j = m$.

This contradicts the optimality of λ^* if $\Delta Z < 0$, or implies that λ' is also an optimal solution to (LP_λ) if $\Delta Z_\lambda(s^*) = 0$. In the latter case, we compute the change in discrepancy from λ^* to λ' . Notice that $\max\{h_{j-2} - h_{j-1}, 0\}$ (or $\max\{v_{s^*-1} - h_{s^*}, 0\}$ if $j = s^* + 1$) can increase by at most ε/f_{j-1} , and the term $\max\{h_{j-1} - h_j, 0\}$ goes down by $\varepsilon/f_{j-1} + \varepsilon/f_j$. The contribution to discrepancy from other terms is unchanged. Hence the discrepancy changes by $\varepsilon/f_{j-1} - (\varepsilon/f_{j-1} + \varepsilon/f_j) = -\varepsilon/f_j < 0$, contradicting our choice of λ^* as the optimal solution with the smallest discrepancy.

Case 2: $h_{s^*}^* \leq h_{s^*+1}^* \leq \dots \leq h_m^*$ but $v_{s^*-1} > h_{s^*}^*$. Let l be the largest index such that $h_{s^*}^* = h_{s^*+1}^* = \dots = h_l^* < h_{l+1}^* + 1$.

We construct a contradiction by considering a new set of $\{\lambda'_i\}_{i=0}^m$, such that

$$\begin{aligned}\lambda'_i &= \lambda_i^* - \varepsilon \quad \forall i \in [s^*, l], \\ \lambda'_i &= \lambda_i^* \quad \forall i \in [1, j-2] \cup [l+1, m].\end{aligned}$$

Denote $h'_i = i - \frac{1-F(i)}{f_i} - \frac{\lambda'_i}{f_i} + \frac{\lambda'_{i-1}}{f_i}$ for all i . This change results in the following changes to the values of $\{h_i^*\}_{i=0}^m$:

$$\begin{aligned}h'_{s^*} &= h_{s^*}^* + \frac{\varepsilon}{f_j}, \\ h'_{l+1} &= h_{l+1}^* - \frac{\varepsilon}{f_{l+1}}, \\ h'_i &= h_i^* \quad \forall i \notin \{s^*, l+1\}.\end{aligned}$$

Denote the change in the (LP_λ) problem objective function from changing λ^* to λ' by $\Delta Z_\lambda(s^*)$. Consider the pairs (t, s) , for which $W(t, s)$ are affected by this change.

For $\varepsilon > 0$ sufficiently small, increasing $h_{s^*}^*$ by $\frac{\varepsilon}{f_{s^*}}$ affects $W(t, s)$ only if $v_t \leq h_{s^*}^*$. Similarly, the decrease of h_{l+1}^* by $\frac{\varepsilon}{f_{l+1}}$ affects $W(t, s)$ if $v_t < h_{l+1}^*$.

Therefore the change in the objective function $\Delta Z_\lambda(s^*)$ is:

$$\Delta Z_\lambda(s^*) \leq \varepsilon \left[\sum \{f_t : v_t \leq h_{s^*}^*\} \right] - \varepsilon \left[\sum \{f_t : v_t < h_{l+1}^*\} \right].$$

Hence $\Delta Z_\lambda(s^*) \leq 0$, and we conclude that λ' is also an optimal solution to (LP_λ) .

Computing the change in discrepancy from λ^* to λ' we observe that $[v_{s^*-1} - h_{s^*}^*]$ will decrease by ε/f_{s^*} . The term $\max\{h_{s^*}^* - h_{s^*+1}^*, 0\}$ goes up

by ε/f_{s^*} . The term $\max\{h_l^* - h_{l+1}^*, 0\}$ goes down by ε/f_{l+1} . The contribution to discrepancy from other terms is unchanged.

Notice that the discrepancy changes by $\varepsilon/f_{s^*} - \varepsilon/f_{s^*} - \varepsilon/f_{l+1} < 0$, contradicting our choice of λ^* as the one with the smallest discrepancy. ■

Lemma 4 *There is an optimal solution (W^*, λ^*) to (LP_λ) such that for any $j \geq s^*$*

$$\{t : v_t \leq h_j^*\} = \{t : v_t < h_{j+1}^*\}.$$

Proof

Let (W^*, λ^*) be the optimal solution to (LP_λ) identified in Lemma 3. If the Lemma is false there is a $j \geq s^*$ such that $h_j^* < h_{j+1}^*$. We construct a contradiction by considering a new set of $\{\lambda'_i\}_{i=0}^m$, such that

$$\begin{aligned}\lambda'_j &= \lambda_j^* - \varepsilon, \\ \lambda'_i &= \lambda_i^* \quad \forall i \neq j\end{aligned}$$

Denote $h'_i = i - \frac{1-F(i)}{f_i} - \frac{\lambda'_i}{f_i} + \frac{\lambda'_{i-1}}{f_i}$ for all i . This change results in the following changes to the values of $\{h_i^*\}_{i=0}^m$:

$$\begin{aligned}h'_j &= h_j^* + \frac{\varepsilon}{f_j}, \\ h'_{j+1} &= h_{j+1}^* - \frac{\varepsilon}{f_{j+1}}, \\ h'_i &= h_i^* \quad \forall i \notin \{j, j+1\}.\end{aligned}$$

Denote the change in the (LP_λ) problem objective function from changing λ^* to λ' by $\Delta Z_\lambda(s^*)$. Consider the pairs (t, s) , for which $W(t, s)$ are affected by this change.

For $\varepsilon > 0$ sufficiently small, increasing h_j^* by $\frac{\varepsilon}{f_j}$ affects $W(t, s)$ only if $v_t \leq h_j^*$. Similarly, the decrease of h_{j+1}^* by $\frac{\varepsilon}{f_{j+1}}$ affects $W(t, s)$ if $v_t < h_{j+1}^*$.

Therefore the change in the objective function $\Delta Z_\lambda(s^*)$ is:

$$\Delta Z_\lambda(s^*) \leq \varepsilon[\sum\{f_t : v_t \leq h_j^*\}] - \varepsilon[\sum\{f_t : v_t < h_{j+1}^*\}].$$

Since $h_j^* < h_{j+1}^*$ the set $\{t : v_t \leq h_j^*\}$ is a subset of $\{t : v_t < h_{j+1}^*\}$. If it is a strict subset this would mean $\Delta Z_\lambda(s^*) < 0$ contradicting optimality of (W^*, λ^*) . ■

Theorem 6 Let (W^*, λ^*) be the optimal solution identified in Lemma 4. There is an optimal solution (a^*, a_b^*) to (P_{λ^*}) such that

1. $a^*(t, s) \leq a^*(t + 1, s)$ for all t ,
2. $a_b^*(t, s) \leq a_b^*(t, s + 1)$ for all s , and
3. $a_b^*(t, s^*) = a_b^*(t, s^* + 1) = \dots = a_b^*(t, m)$.

Proof

Now

$$Z_{\lambda^*}(s^*) = \max \sum_{t=1}^m \sum_{s=1}^m f_t v_t f_s a(t, s) + \sum_{s=1}^{s^*-1} \sum_{t=1}^m f_s v_s f_t a_b(t, s) + \sum_{s=s^*}^m \sum_{t=1}^m f_s h_s^* f_t a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1 \quad \forall t, s$
 $a(t, s), a_b(t, s) \geq 0 \quad \forall s, t$

This decomposes into a collection of subproblems, one subproblem for each profile (t, s) of types. When $s \leq s^* - 1$ the subproblem is

$$g_{\lambda^*}(t, s) = \max v_t a(t, s) + v_s a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1$
 $a(t, s), a_b(t, s) \geq 0$

and when $s \geq s^*$ it is

$$g_{\lambda^*}(t, s) = \max v_t a(t, s) + h_s^* a_b(t, s)$$

s.t. $a(t, s) + a_b(t, s) \leq 1$
 $a(t, s), a_b(t, s) \geq 0$

For the case when $s \leq s^* - 1$ the optimal solution is to award the good to the agent with highest non-negative virtual value. We will break ties in favor of the unconstrained bidder. Specifically, when $s \leq s^* - 1$ we have the following.

1. When $v_t \geq \max\{v_s, 0\} \Rightarrow a^*(t, s) = 1$.
2. When $v_s > \max\{v_t, 0\} \Rightarrow a_b^*(t, s) = 1$.

When $s \geq s^*$ we have the following.

1. When $v_t \geq \max\{h_s^*, 0\} \Rightarrow a^*(t, s) = 1$.
2. When $h_s^* > \max\{v_t, 0\} \Rightarrow a_b^*(t, s) = 1$.

Monotonicity of the virtual values and the h^* 's (from Lemma 3) yield items 1 and 2 of the Theorem.

To prove item 3, suppose it is false. Then there is a $j \geq s^*$ and t such that

$$a_b^*(j+1, t) = 1 \neq 0 = a_b^*(j, t).$$

This can happen only if $h_j^* \leq v_t < h_{j+1}^*$. However, this contradicts Lemma 4, since $\{t : v_t \leq h_j^*\} = \{t : v_t < h_{j+1}^*\}$. \blacksquare

3.2 Solving $[BOPT(s^*)]$

The optimal solution to (P_{λ^*}) satisfies the omitted monotonicity constraints and is therefore an optimal solution to $[BOPT(s^*)]$. In particular, Theorem 6 establishes there is an optimal solution to $[BOPT(s^*)]$ such that $a_b(t, s) = a_b(t, s^*)$ for all t and all $s \geq s^*$.

This observation allows us to rewrite $[BOPT(s^*)]$ as

$$Z(s^*) = \max_{s,t} \sum [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s]$$

$$\text{s.t. } \mathcal{A}_1^b \leq \dots \leq \mathcal{A}_m^b$$

$$\mathcal{A}_1 \leq \dots \leq \mathcal{A}_m$$

$$\mathcal{A}_i = \sum_s f_s a(i, s) \quad \forall i$$

$$\mathcal{A}_i^b = \sum_t f_t a_b(t, i) \quad \forall i$$

$$a_b(t, s^*) = \dots = a_b(t, m) \quad \forall t$$

$$a(t, s) + a_b(t, s) \leq 1 \quad \forall t, s$$

$$a(t, s), a_b(t, s) \geq 0 \quad \forall t, s$$

We will solve this version of $[BOPT(s^*)]$ by removing the monotonicity constraints and arguing that the solution to the relaxed problem satisfies them.

$$\begin{aligned}
F(s^*) &= \max \sum_{s,t} [f_t \mathcal{A}_t v_t + f_s \mathcal{A}_s^b v_s] \\
\text{s.t. } \mathcal{A}_i &= \sum_s f_s a(i, s) \quad \forall i \\
\mathcal{A}_i^b &= \sum_t f_t a_b(t, i) \quad \forall i \\
a_b(t, s^*) &= \dots = a_b(t, m) \quad \forall t \\
a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\
a(t, s), a_b(t, s) &\geq 0 \quad \forall t, s
\end{aligned}$$

This last program is equivalent to

$$\begin{aligned}
F(s^*) &= \max \sum_{t=1}^m \sum_{s=1}^m f_t f_s a(t, s) v_t + \sum_{t=1}^m \sum_{s=1}^m f_t f_s a_b(t, s) v_s \\
\text{s.t. } a_b(t, s^*) &= \dots = a_b(t, m) \quad \forall t \\
a(t, s) + a_b(t, s) &\leq 1 \quad \forall t, s \\
a(t, s), a_b(t, s) &\geq 0 \quad \forall t, s
\end{aligned}$$

Eliminating the $a_b(t, s)$ variables for $s \geq s^* + 1$ we can rewrite the program as

$$\begin{aligned}
F(s^*) &= \max \sum_{t=1}^m \sum_{s=1}^m f_t f_s a(t, s) v_t + \sum_{t=1}^m \sum_{s=1}^{s^*-1} f_t f_s a_b(t, s) v_s + \sum_{t=1}^m f_t \left[\sum_{j=s^*}^m f_j v_j \right] a_b(t, s^*) \\
\text{s.t. } a(t, s) + a_b(t, s) &\leq 1 \quad \forall t = 1, \dots, m \quad \forall s = 1, \dots, s^* \\
a, a_b &\geq 0
\end{aligned}$$

Set $w_s = v_s$ for $s \leq s^* - 1$ and $w_{s^*} = \frac{\sum_{j=s^*}^m f_j v_j}{\sum_{j=s^*}^m f_j}$. Monotonicity of the v_s 's implies that the w_s 's are monotone as well. Set $g_j = f_j$ for $j \leq s^* - 1$ and $g_{s^*} = \sum_{j=s^*}^m f_j$. Then

$$F(s^*) = \max \sum_{t=1}^m \sum_{s=1}^{s^*} f_t g_s a(t, s) v_t + \sum_{t=1}^m \sum_{s=1}^{s^*} f_t g_s a_b(t, s) w_s$$

$$\begin{aligned} \text{s.t. } a(t, s) + a_b(t, s) &\leq 1 \quad \forall t = 1, \dots, m \quad \forall s = 1, \dots, s^* \\ a(t, s), a_b(t, s) &\geq 0 \quad \forall t = 1, \dots, m \quad \forall s = 1, \dots, s^* \end{aligned}$$

This decomposes into a collection of subproblems one for each profile (t, s) :

$$\begin{aligned} \max v_t a(t, s) + w_s a_b(t, s) \\ \text{s.t. } a(t, s) + a_b(t, s) &\leq 1 \\ a(t, s), a_b(t, s) &\geq 0 \end{aligned}$$

Therefore, if we know the threshold, s^* , the optimal mechanism could be described thus:

1. If $v_t > \max\{w_s, 0\}$ then $a(t, s) = 1$ and $a_b(t, s) = 0$.
2. If $w_s \geq \max\{v_t, 0\}$ then $a_b(t, s) = 1$ and $a(t, s) = 0$.
3. If $v_t, w_s < 0$ set $a_b(t, s) = a(t, s) = 0$.

Monotonicity of the v 's and w 's ensures that the omitted monotonicity constraints are satisfied.

To determine the payments we use the fact that

$$p_s^b = \sum_{j=0}^{s-1} (j+1)(\mathcal{A}_{j+1}^b - \mathcal{A}_j^b) = s\mathcal{A}_s^b - \sum_{j=1}^{s-1} \mathcal{A}_j^b$$

and

$$p_t = \sum_{j=0}^{t-1} (j+1)(\mathcal{A}_{j+1} - \mathcal{A}_j) = t\mathcal{A}_t - \sum_{j=1}^{t-1} \mathcal{A}_j$$

for all s, t .

Let r_f be the smallest index such that $v_{r_f} \geq 0$ and $r(s^*)$ the smallest index less than s^* such that $w_{r(s^*)} \geq 0$. Then $\mathcal{A}_t = 0$ when $t < r_f$ and $\mathcal{A}_t = \sum\{f_s : w_s < v_t\} = Pr(w_s < v_t)$ when $t \geq r_f$. Similarly, $\mathcal{A}_s^b = 0$ when $s < r(s^*)$ and $\mathcal{A}_s^b = \sum\{f_t : v_t \leq w_s\} = Pr(v_t \leq w_s)$. Therefore, $p_s^b = 0$ for $s < r(s^*)$ and

$$p_s^b = sPr(v_t \leq w_s) - \sum_{j=1}^{s-1} Pr(v_t \leq w_j) \quad \forall s \geq r(s^*).$$

Similarly, $p_t = 0$ for $t < r_f$ and

$$p_t = tPr(w_s < v_t) - \sum_{j=1}^{t-1} Pr(w_s < v_j) \quad \forall t \geq r_f.$$

3.2.1 Determining s^*

A condition that any candidate for the threshold s^* must satisfy is that

$$b \geq p_{s^*}^b = s^* Pr(v_t \leq w_{s^*}) - \sum_{j=1}^{s^*-1} Pr(v_t \leq w_j).$$

For $j \leq s^* - 1$ we know that $w_j = v_j$. Therefore s^* can be chosen to be the largest index s such that

$$s Pr(v_t \leq \frac{\sum_{i=s}^m f_i v_i}{\sum_{i=s}^m f_i}) - \sum_{j=1}^{s-1} Pr(v_t \leq v_j) \leq b.$$

3.3 A Description of the Optimal Mechanism

The optimal mechanism can be described in a way that is very similar to the description of the optimal mechanism in Myerson (1981).

1. First determine the optimal threshold s^* .
2. Compute for each type s of the constrained bidder a *modified* virtual value w_s as follows. If $s \leq s^* - 1$ set $w_s = v_s$ and if $s \geq s^*$ set $w_s = \frac{\sum_{j=s^*}^m f_j v_j}{\sum_{j=s^*}^m f_j}$.
3. Compute for each type t of the unconstrained bidder its virtual value.
4. If the unconstrained types virtual value exceeds the constrained types modified virtual value and is non-negative award the good to the unconstrained bidder. If the modified virtual value of the constrained bidder is at least as large as the virtual value of the unconstrained virtual value and is non-negative, award the good to the constrained bidder.

For this reason we will call the mechanism the threshold mechanism.

4 Private Budget Constraint

We turn now to the case when the budget of the constrained bidder is private information as well. In our mechanism the constrained bidder must post a bond equal to their reported budget. This ensures that the constrained

bidders payment cannot exceed their reported budget. In terms of incentive compatability this will mean that we can ignore the possibility that the constrained bidder will inflate her budget. This allows us to focus on the case of a constrained bidder with a large budget pretending to have a smaller budget.

We discuss the Bayesian incentive compatible case only. Our goal will be to reduce the problem to the one considered in section 3 of this paper. A similar analysis applies in the dominant strategy incentive compatible case.

Let \mathcal{A}_{ib} be the expected allocation to the constrained agent when the constrained agent reports (i, b) . Her expected payment will be p_{ib} . The requirement that the constrained bidder must post a bond implies that the constrained bidder can only underreport her budget. Therefore we only have to consider downward incentive constraints with respect to the reported budget, i.e.

$$i\mathcal{A}_{ib} - p_{ib} \geq i\mathcal{A}_{ib'} - p_{ib'} \quad \forall b' < b. \text{ (DIC)}$$

We argue that the optimal mechanism in the case of a private budget constraint under the bond requirement is implemented through a variation of the threshold mechanism derived in section 3 of this paper. First, the bidders post bonds, revealing their budget constraints,. Then the threshold mechanism is implemented based on the revealed information about b . To prove optimality of the procedure it suffices to show that it satisfies (DIC).

Theorem 7 *The optimal mechanism when the constrained bidder must post a bond is:*

$$\mathcal{A}_{ib} = \mathcal{A}_i^b, p_{ib} = p_i^b.$$

Proof

It suffices to verify that (DIC) is satisfied. Let s_b, s_c be the threshold associated with budgets b and c where $b > c$. It is easy to see that $s_b \geq s_c$. If the constrained type is i , denote by w_i^b and w_i^c the modified virtual values associated with budgets b and c respectively. We have three cases.

1. $i < s_c$

Since $w_i^b = w_i^c$,

$$\mathcal{A}_{ib} = \mathcal{A}_i^b = Pr(v_t \leq w_i^b) = Pr(v_t \leq w_i^c) = \mathcal{A}_i^c = \mathcal{A}_{ic}.$$

A similar argument shows that $p_{ib} = p_i^b = p_i^c = p_{ic}$. Therefore (DIC) holds.

2. $s_c \leq i < s_b$

As the threshold mechanism is incentive compatible with respect to reports of the value $i\mathcal{A}_{ib} - p_{ib} = i\mathcal{A}_i^b - p_i^b \geq i\mathcal{A}_{s_c}^b - p_{s_c}^b$. Now

$$i\mathcal{A}_{s_c}^b - p_{s_c}^b = iPr(v_t \leq w_{s_c}^b) - [s_c Pr(v_t \leq w_{s_c}^b) - \sum_{j=1}^{s_c-1} Pr(v_r \leq w_j^b)].$$

Since $w_j^b = v_j$ for all $j < s_b$ and $w_j^b = w_j^c = v_j$ for all $j < s_c$ we have that

$$i\mathcal{A}_{s_c}^b - p_{s_c}^b = iPr(v_t \leq v_{s_c}) - [s_c Pr(v_t \leq v_{s_c}) - \sum_{j=1}^{s_c-1} Pr(v_t \leq w_j^c)].$$

Since $v_{s_c} \leq w_{s_c}^c$, the term on the right hand side is

$$\begin{aligned} & iPr(v_t \leq v_{s_c}) + iPr(v_{s_c} < v_t \leq w_{s_c}^c) - [iPr(v_{s_c} < v_t \leq w_{s_c}^c) + s_c Pr(v_t \leq v_{s_c}) - \sum_{j=1}^{s_c-1} Pr(v_t \leq w_j^c)] \\ & \geq iPr(v_t \leq w_{s_c}^c) - [iPr(v_t \leq v_{s_c}) - \sum_{j=1}^{s_c-1} Pr(v_t \leq w_j^c)] = i\mathcal{A}_i^c - p_i^c = i\mathcal{A}_{ic} - p_{ic}. \end{aligned}$$

3. $i \geq s_b$

Incentive compatibility in the valuations and monotonicity of \mathcal{A}^b implies

$$i\mathcal{A}_{ib} - p_{ib} = i\mathcal{A}_{s_b}^b - p_{s_b}^b \geq i\mathcal{A}_{s_c}^b - p_{s_c}^b.$$

Now

$$\begin{aligned} i\mathcal{A}_{s_c}^b - p_{s_c}^b &= iPr(v_t \leq v_{s_c}) - [iPr(v_t \leq v_{s_c}) - \sum_{j=1}^{s_c-1} Pr(v_t \leq v_j)] \\ &= iPr(v_t \leq w_{s_c}^c) - [iPr(v_t \leq w_{s_c}^c) - \sum_{j=1}^{s_c-1} Pr(v_t \leq v_j)] = i\mathcal{A}_i^c - p_i^c. \end{aligned}$$

■

5 Acknowledgements

We thank Peter Eso, Peter Klibanoff, Rudolf Muller and Hongwei Gui for useful comments.

6 References

1. Che, Y.-K., and I. Gale, 1996, "Expected Revenue of All-Pay Auctions and First-Price Sealed-Bid Auctions with Budget Constraints," *Economics Letters*, 50, 373-379.
2. Che, Y.-K., and I. Gale, 2000, "The Optimal Mechanism for Selling to a Budget-Constrained Buyer," *Journal of Economic Theory*, 92, 198-233.
3. Gavious, A. B. Moldovanu and A. Sela, 2002, "Bid Costs and Endogenous Bid Caps," *RAND Journal of Economics*, vol. 33(4), pages 709-722
4. Harris, M. and A. Raviv (1981), "A Theory of Monopoly Pricing Schemes with Demand Uncertainty," *American Economic Review*, 71, pp. 347-65.
5. Laffont, J.-J., and J. Robert, 1996, "Optimal Auction with Financially Constrained Buyers," *Economics Letters*, 52, 181-186.
6. Maskin, E. S., 2000, "Auctions, Development, and Privatization: Efficient Auctions with Liquidity- Constrained Buyers," *European Economic Review*, 44, 667-681.
7. Myerson, R., 1981, "Optimal Auction Design," *Mathematics of Operations Research*, 6, 58-73.
8. Zheng, C. Z., 2001, "High Bids and Broke Winners," *Journal of Economic Theory*, 100, 129-171.