

Discussion Paper No. 751

THE POLYTOPE OF DEGREE SEQUENCES

by

Uri N. Peled*

and

Murali K. Srinivasan**

November 1987

*Department of Managerial Economics and Decision Sciences, J.L. Kellogg Graduate School of Management, Northwestern University, Evanston, Illinois 60208. (Electronic mail: U32799@UICVM.BITNET.)

**Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, Box 4348, Chicago, Illinois 60680. (Electronic mail: U13720@UICVM.BITNET.)

Abstract

A nonnegative integer sequence (d_1, d_2, \dots, d_n) is called a degree sequence if there exists a simple graph on the vertex set $V = \{1, 2, \dots, n\}$ such that $\deg(i) = d_i$, for all i . The degree sequence of a threshold graph is a threshold sequence. Let $D_n = \text{Convex Hull} \{(x_1, x_2, \dots, x_n) \mid (x_1, \dots, x_n) \text{ is a degree sequence}\}$. It is proved that:

- 1) A degree sequence f is an extreme point of D_n if and only if f is a threshold sequence.
- 2) Two threshold sequences f and g are adjacent extreme points of D_n if and only if f can be obtained from g by either adding 1 to two components of g or subtracting 1 from two components of g .
- 3) D_n is determined by the following system of inequalities:

$$\sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$$

for all set S, T with $\emptyset \neq S \cup T \subseteq \{1, 2, \dots, n\}$, $S \cap T = \emptyset$

Moreover, this system is totally dual integral. Furthermore, for $n \geq 4$, (S, T) determines a facet of D_n if and only if either $|S \cup T| = 1$ or else $S \neq \emptyset$, $T \neq \emptyset$, $|S \cup T| \neq n - 1, n - 2$.

- 4) f is a threshold sequence if and only if the only degree sequences majorizing f in the sense of Hardy-Littlewood-Pólya are the rearrangements of f .
- 5) Every degree sequence is a convex combination of isomorphic threshold sequences (i.e., threshold sequences that are rearrangements of each other).

i. Introduction

A nonnegative integer sequence $d = (d_1, d_2, \dots, d_n)$ is called a degree sequence if there exists a (simple) graph $G = (V, E)$ on the vertex set $V = \{1, 2, \dots, n\}$ such that $\deg(i) = d_i$, for all i . G is said to be a realization of d . Degree sequences have been thoroughly studied and the following are three well-known characterizations of nonincreasing degree sequences ($d_1 \geq d_2 \geq \dots \geq d_n$):

- (1) $\sum_{i=1}^n d_i$ is even and (d_1, \dots, d_n) satisfies the Erdős-Gallai inequalities $\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k)$, $k = 1, \dots, n$ [2,5,16];
- (2) $\sum_{i=1}^n d_i$ is even and (d_1, \dots, d_n) is majorized (Definition 5.1 below) by its corrected conjugate sequence $(\bar{d}_1, \dots, \bar{d}_n)$ given by

$$\bar{d}_k = |\{i < k | d_i \geq k-1\}| + |\{i > k | d_i \geq k\}| \quad [2];$$

- (3) $(d_2 - 1, d_3 - 1, \dots, d_{d_1+1} - 1, d_{d_1+2}, \dots, d_n)$ is also a degree sequence.

This criterion, when iterated, gives rise to the Havel-Hakimi algorithm [8,13].

Let $DS_n = \{(x_1, \dots, x_n) | (x_1, \dots, x_n) \text{ is a degree sequence}\}$. The polytope of degree sequences is $D_n = \text{Convex Hull}(DS_n)$. In [15], Koren proved that:

- (1) A degree sequence is an extreme point of D_n if and only if it has a unique realization (see also [9]);
- (2) D_n is determined by the following system of linear inequalities:

$$\sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$$

for all set S, T such that $\emptyset \neq S \cup T \subseteq \{1, 2, \dots, n\}$, $S \cap T = \emptyset$.

Koren also characterized the uniquely realizable nonincreasing degree sequences by the property $d = \bar{d}$. The proofs in [15] rely on the Erdős-Gallai inequalities.

In this paper we study the properties of D_n using threshold graphs. In Sections 2 and 4 we reproduce the results of [15] using linear programming duality and the structure of threshold graphs and show that the above system of linear inequalities is totally dual integral. This approach fits naturally the modern trend of using LP duality and some combinatorics to obtain polyhedral results [20]. We also determine the facets of D_n in Section 4. In Section 3 we study the problem of adjacency of extreme points of D_n and show that two extreme points (f_1, \dots, f_n) and (g_1, \dots, g_n) of D_n with realizations $G_1 = (V, E_1)$ and $G_2 = (V, E_2)$ (which are necessarily unique) are adjacent if and only if $|E_1 \ominus E_2| = 1$ (\ominus denotes the symmetric difference).

Finally, in Section 5 we show that the non-majorizable degree sequences (more precisely, degree sequences d that cannot be majorized by other degree sequences except for rearrangements of d) are precisely the degree sequences of threshold graphs. From a well-known result in the theory of majorization it then follows that every degree sequence is a convex combination of degree sequences of isomorphic threshold graphs.

The following definitions are used in this paper. All graphs considered are finite, undirected and without loops or parallel edges. $[a, b]$ denotes an edge joining two vertices a and b , and $N(a) = \{b \mid [a, b] \text{ is}$

an edge} is the set of neighbors of a . $N(W) = \cup \{N(a) | a \in W\}$ for $W \subseteq V$. A clique is a set of pairwise adjacent vertices and a stable set is a set of pairwise nonadjacent vertices. The size of the largest clique of a graph $G = (V, E)$ is written as $\omega(G)$. The vertex set of each graph discussed here is assumed to be $\{1, 2, \dots, n\}$, for some n . We denote by u_i the i -th unit vector.

Threshold graphs were introduced by Chvátal and Hammer [3] as a class of graphs for which there is a simple method of distinguishing stable sets from nonstable sets. A graph with vertex set $\{1, 2, \dots, n\}$ is a threshold graph if there exist real weights w_1, w_2, \dots, w_n , t such that the 0-1 solutions of the inequality $\sum_{i=1}^n w_i x_i \leq t$ are precisely the characteristic vectors of the stable sets of the graph. The degree sequence of a threshold graph is called a threshold sequence.

Threshold graphs have many characterizations. We present the two that are used in the sequel.

Theorem 1.1 [3]: A graph is threshold if and only if it has no induced subgraphs isomorphic to $2K_2$, P_3 or C_4 , as shown in Figure 1. \square

INSERT FIGURE 1 ABOUT HERE

Equivalently, Theorem 1.1 could be stated as:

Corollary 1.2: A graph is threshold if and only if it does not contain the configuration shown in Figure 2, where solid lines represent edges, dotted lines represent non-edges, and the absence of a line allows for the

possibility of the edge existing or not. \square

INSERT FIGURE 2 ABOUT HERE

Theorem 1.3 [3]: A graph $G = (V, E)$ is threshold if and only if there is a partition of V into disjoint sets K and I (possibly empty) such that

- (1) K is a clique;
- (2) I is a stable set;
- (3) There exists an ordering u_1, u_2, \dots, u_s of the vertices of I such that

$$N(u_1) \subseteq N(u_2) \subseteq \dots \subseteq N(u_s)$$

or equivalently, there exists an ordering v_1, v_2, \dots, v_k of the vertices of K such that

$$N(v_1) \cap I \subseteq N(v_2) \cap I \subseteq \dots \subseteq N(v_k) \cap I. \quad \square$$

It is shown in [7] (see also [10]) that we can always choose K with $|K| = \omega(G)$. A partition satisfying (1) and (2) of Theorem 1.3 is called a split partition.

Theorem 1.4 [7] (see also [9]): A nonnegative integer sequence d is a threshold sequence if and only if $d = \bar{d}$. \square

In light of Theorem 1.4, Koren's results about the extreme points of D_n can be interpreted to mean that the extreme points of D_n are precisely the

threshold sequences. For further characterizations and properties of threshold graphs and threshold sequences see [1, 3, 6, 7, 9, 11, 14, 18, 19].

2. Extreme Points

We first review a few facts about the structure of threshold graphs.

Lemma 2.1: Let $G = (V, E)$ be a threshold graph with split partition $V = K \cup I$, $|K| = \omega(G)$. Then there exists a $v \in K$ that is not adjacent to any vertex in I .

Proof: Enumerate the vertices of K as v_1, v_2, \dots, v_k so that

$$N(v_1) \cap I \subseteq N(v_2) \cap I \subseteq \dots \subseteq N(v_k) \cap I.$$

Then v_1 is not adjacent to any vertex in I . For if $u \in N(v_1) \cap I$, then $K \cup u$ is a larger clique, contradicting our choice of K . \square

We now introduce a partition of the vertices of a threshold graph, which is very similar to the degree partition of [7].

Definition 2.2: Let $G = (V, E)$ be a threshold graph with a split partition $V = K \cup I$, $|K| = \omega(G)$.

If the edges in K are omitted, one obtains a bipartite graph $B = (K, I; F)$. Let $k_0 < k_1 < \dots < k_p$ be the distinct degrees in B of the vertices in K . By Lemma 2.1, $k_0 = 0$. Let $s_1 < s_2 < \dots < s_p$ be the

distinct nonzero degrees in B of the vertices of I (by Theorem 1.3, the number of distinct positive degrees in I is p). Set $s_0 = 0$, and define for $i = 0, 1, \dots, p$,

$$K_i = \{v \in K \mid \deg_B(v) = k_i\}$$

$$I_i = \{v \in K \mid \deg_B(v) = s_i\}$$

By this definition all the K_i and I_i are nonempty, with the possible exception of I_0 . See Figure 3.

It follows from Definition 2.2 that

$$N(K_i) \cap I = I_{p-i+1} \cup \dots \cup I_p$$

$$N(I_i) = K_{p-i+1} \cup \dots \cup K_p$$

for $i = 0, 1, \dots, p$.

Remark: The fact that $K_0 = \{v \in K \mid N(v) \cap I = \emptyset\} \neq \emptyset$ was derived from the choice of K to satisfy $|K| = \omega(G)$. It is needed in some, but not all of the proofs below. We assume it in all cases for uniformity.

INSERT FIGURE 3 ABOUT HERE

Let $(c_1, c_2, \dots, c_n) \in \mathbb{R}^n$. Consider the combinatorial optimization problem

$$\max \sum_{i=1}^n c_i x_i$$

s.t. $(x_1, x_2, \dots, x_n) \in D_n$
 (or equivalently, $(x_1, x_2, \dots, x_n) \in DS_n$).

The following very simple algorithm produces an optimal solution:

Algorithm 2.3:

$V \leftarrow \{1, 2, \dots, n\};$

$E \leftarrow \emptyset;$

for all $i, j, i \neq j$, if $c_i - c_j \geq 0$, add edge $[i, j]$ to E ;

$(x_1, x_2, \dots, x_n) \leftarrow$ degree sequence of $G = (V, E)$.

Remarks:

2.4 It is clear that the algorithm is correct: adding the edge $[i, j]$ adds $c_i + c_j$ to the objective function.

2.5 If $c_i + c_j = 0$ for some $i, j, i \neq j$, we have the freedom of not adding the edge $[i, j]$ and still obtain an optimal solution. This freedom is enough to obtain all graphs with optimal degree sequences.

2.6 In particular, the following conditions are equivalent:

- (a) $c_i + c_j \neq 0$ for all $i, j, i \neq j$;
- (b) There is only one optimal degree sequence;

(c) There is only one graph with an optimal degree sequence.

2.7 The algorithm can be thought of as the greedy algorithm on the uniform matroid on the edges with edge costs $c_i + c_j$. The algorithm can also be interpreted as the "greedy algorithm" described in [22, Section 19.2] applied to D_n . In [22] it is proved that the "greedy algorithm" on a polymatroid works for all $c \in \mathbb{R}^n$. Here it works on D_n , which is not a polymatroid.

Theorem 2.8: A degree sequence is an extreme point of D_n if and only if it is a threshold sequence.

Proof: ("Only if"): Every extreme point of D_n is the unique optimum for some objective function $\sum_{i=1}^n c_i x_i$, so Algorithm 2.3 must produce it on input (c_1, \dots, c_n) . But the algorithm always produces threshold graphs. Indeed, write $V = K \cup I$ where $K = \{i | c_i \geq 0\}$, $I = \{i | c_i < 0\}$. Then K and I satisfy the condition of Theorem 1.3.

("If"): Let $d = (d_1, d_2, \dots, d_n)$ be a threshold sequence and $G = (V, E)$ be a realization of d . Let K_j, I_j , $0 \leq j \leq p$ be as in Definition 2.2.

Choose $c = (c_1, c_2, \dots, c_n)$ as follows (see Figure 4):

$$c_i = \begin{cases} 2j + 1 & i \in K_j \\ -2p - 2 + 2j & i \in I_j \end{cases}$$

INSERT FIGURE 4 ABOUT HERE

It is easy to check that Algorithm 2.3 with input (c_1, c_2, \dots, c_n)

outputs precisely the graph $G = (V, E)$ and its degree sequence $d = (d_1, \dots, d_n)$. Moreover, by Remark 2.6, d is the only optimal sequence. Therefore d is an extreme point of D_n . \square

Corollary 2.9 [8, 13]: A degree sequence has a unique realization if and only if it is a threshold sequence.

Proof: The "if" part follows from Theorem 2.4 and Remark 2.6. For the "only if" part, let G be a non-threshold graph. Then by Corollary 1.2, G has a forbidden configuration (see Figure 5), and an interchange gives a different graph with the same degree sequence. \square

INSERT FIGURE 5 ABOUT HERE

Remarks:

- 2.10 The "if" part also follows easily from the forbidden configuration characterization of threshold graphs and from the fact that all realizations of a degree sequence can be obtained from each other by interchanges along such configurations (see Berge [2], Chapter 8, Theorem 5).
- 2.11 A threshold sequence can be recognized and its realization constructed in time $O(n \log n)$ [3].
- 2.12 The number of extreme points of D_n (i.e., the number of labelled n -vertex threshold graphs) has been determined in [1].

3. Adjacency

In this section we develop criteria for two extreme points of D_n to be adjacent and produce a formula for the number of adjacent threshold sequences to a given threshold sequence.

Lemma 3.1: Let $T_1 = (V, E_1)$ and $T_2 = (V, E_2)$ be two threshold graphs with degree sequences f and g , respectively. If $|E_1 \ominus E_2| \geq 2$, then f and g are not adjacent extreme points of D_n .

Proof: Assume that f and g are adjacent extreme points of D_n . Then there exists a $c = (c_1, c_2, \dots, c_n)$ such that $\sum_{i=1}^n c_i x_i$ is maximized over DS_n only by f and g .

Choose $e \in E_1 \ominus E_2$. Without loss of generality $e \in E_1$. Consider $G_3 = (V, E_2 \cup e)$ with degree sequence $d = (d_1, d_2, \dots, d_n)$. We have $G_3 \neq T_2$ because $e \in E_2 \cup e$ and $e \notin E_2$. Also $G_3 \neq T_1$, as otherwise $E_1 = E_2 \cup e$ and $E_1 \ominus E_2 = \{e\}$, contradicting $|E_1 \ominus E_2| \geq 2$.

Let $e = [i, j]$. Since $e \in E_1$ and f is an optimal degree sequence, $c_i + c_j \geq 0$ by Algorithm 2.3. Hence $\sum_{i=1}^n c_i d_i \geq \sum_{i=1}^n c_i g_i$. As g is optimal, d is optimal too. Moreover, $d \neq f, g$ since the threshold sequences f, g have the unique realizations T_1 and T_2 , respectively. This contradicts the fact that f and g are the only optimal degree sequences for c , proving the lemma. \square

Lemma 3.2: Let $G = (V, E)$ be a threshold graph with a split partition $V = K \cup I$ as in Definition 2.2.

- (a) Let $x, y \in I$. Then $(V, E \cup [x, y])$ is a threshold graph if and only if $x, y \in I_p$ and $|K_0| = 1$ (i.e., $\deg x = \deg y = |K| - 1$).
- (b) Let $x, y \in K$. Then $(V, E - [x, y])$ is a threshold graph if and only if

$$(x, y \in K_0), \quad \text{if } |K_0| \geq 2$$

$$(x \in K_0, y \in K_1) \text{ or } (x \in K_1, y \in K_0), \text{ if } |K_0| = 1.$$

Proof:

(a) ("Only if"): First assume that $x \in I_i, y \in I_j$ and $i < j$. Since K_0 is nonempty, there exists a $z \in K$ with $[z, y] \notin E$. As x has more non-neighbors than y , there exists a $w \in K, z \neq w$ such that $[x, w] \notin E$ (see Figure 6). Adding edge $[x, y]$ closes a forbidden configuration.

INSERT FIGURE 6 ABOUT HERE

Therefore, $x, y \in I_i$, for some i . Now assume that $i < p$. Vertices x and y have the same degree and so, by Theorem 1.3, they have the same neighbors in K . By our assumption there exist $w, z \in K, w \neq z$ such that both w and z are non-neighbors of both x and y . We reach the same contradiction as above. Thus $x, y \in I_p$. Finally, $|K_0| = 1$, as assuming $|K_0| \geq 2$ gives rise to the same contradiction once more.

("If"): Let $K_0 = \{z\}$. Replace K by $K' = K - \{z\} \cup \{x, y\}$ and I by $I' = I - \{x, y\} \cup \{z\}$. It is easy to check that $G' = (K' \cup I', E \cup [x, y])$ satisfies the condition of Theorem 1.3, proving that G' is threshold.

(b) ("Only if")

Case 1: $|K_0| \geq 2$. Assume that $x \in K_i$, $i \geq 1$, $y \in K_j$, $j \geq 0$, $i \geq j$.

This means that there exists a $z \in K_0$, $z \neq y$ and a $w \in I$ such that $[x,w] \in E$ (Figure 7). Dropping $[x,y]$ opens up a forbidden configuration. Therefore, $x,y \in K_0$.

INSERT FIGURE 7 ABOUT HERE

Case 2: $|K_0| = 1$. Let $K_0 = \{z\}$. First assume that $x \in K_i$, $y \in K_j$, $i,j \geq 1$. There exists a $w \in I$ with $[x,w] \in E$. We reach the same contradiction as in Case 1.

Now assume without loss of generality that $y \in K_0$ (hence $y = z$) and $x \in K_i$, $i \geq 2$. Choose $z_1 \in K_1$ and $w \in I$ with $[x,w] \in E$ and $[z_1,w] \notin E$ (Figure 8).

INSERT FIGURE 8 ABOUT HERE

Dropping $[x,y]$ opens up a forbidden configuration. Therefore,

$$(x \in K_0, y \in K_1) \text{ or } (x \in K_1, y \in K_0).$$

("If):

Case 1: $|K_0| \geq 2$. Replace K by $K' = K - \{x,y\}$, and I by $I' = I \cup \{x,y\}$.

Case 2: $|K_0| = 1$. Assume without loss of generality that $x \in K_1$, $y \in K_0$. Replace K by $K' = K - \{y\}$, and I by $I' = I \cup \{y\}$. In both cases it

is easy to check that $G = (K' \cup I', E - [x,y])$ satisfies the condition of Theorem 1.3, proving that G' is threshold. \square

Lemma 3.3: Let $G = (V,E)$ be a threshold graph with split partition $V = K \cup I$ as in Definition 2.2.

- (a) Let $x \in I_r$ and $y \in K_s$, $[x,y] \notin E$. Then $(V, E \cup [x,y])$ is threshold if and only if $r + s = p$.
- (b) Let $x \in I_r$ and $y \in K_s$, $[x,y] \in E$. Then $(V, E - [x,y])$ is threshold if and only if $r + s = p + 1$.

Proof:

(a) ("Only if"): Since $[x,y] \notin E$, $r + s \leq p$. Assume that $r + s \leq p - 1$. Pick $z \in I_{r+1}$ and $w \in K_{p-r}$. Then $[x,w] \notin E$, $[z,y] \notin E$ and $[z,w] \in E$ (Figure 9). Adding edge $[x,y]$ closes a forbidden configuration. Hence $r + s = p$.

INSERT FIGURE 9 ABOUT HERE

("If"): One can easily verify that $(K \cup I, E \cup [x,y])$ continues to satisfy the condition of Theorem 1.3.

- (b) Similar to (a). \square

Lemma 3.4: Let $G = (V,E)$ be a threshold graph and $[x,y] \notin E$. If $G' = (V, E \cup [x,y])$ is also a threshold graph, then G has a split partition $V = K \cup I$ such that $(x \in I, y \in K)$ or $(x \in K, y \in I)$, $|K| = \omega(G)$.

Proof: Let $V = K \cup I$ be a split partition of G as in Definition 2.2. If $x, y \in I$ and G' is threshold, then by Lemma 3.2 $x, y \in I_p$ and $|K_0| = 1$. Let $K_0 = \{z\}$. Replacing K by $K - \{z\} \cup \{x\}$ and I by $I - \{x\} \cup \{z\}$ we get another split partition with the required properties. \square

Theorem 3.5: Let $T_1 = (V, E_1)$ and $T_2 = (V, E_2)$ be the (unique) realizations of the threshold sequences f and g , respectively. Then f and g are adjacent extreme points of D_n if and only if $|E_1 \ominus E_2| = 1$.

Proof: The "only if" direction is Lemma 3.1.

("If"): Let $E_1 \ominus E_2 = \{e\}$. Without loss of generality, $E_1 = E_2 \cup \{e\}$. From Lemma 3.4 we can assume that T_2 has a split partition $V = K \cup I$ such that $x \in I$, $y \in K$, where $[x, y] = e$. Let $x \in I_r$ and $y \in K_s$. By Lemma 3.3, $r + s = p$. Run Algorithm 2.3 on the input $c = (c_1, c_2, \dots, c_n)$ defined as follows (see Figure 10):

$$\begin{aligned}
 c_i &= -2p + 2r - 2, & i &= x \\
 c_i &= 2p - 2r + 2, & i &= y \\
 c_i &= -2p + 2j - 2, & i &\in I_j, j \geq r + 1 \\
 c_i &= -2p + 2j - 4, & i &\in I_j, i \neq x, j \leq r \\
 c_i &= 2j + 1, & i &\in K_j, i \neq y, j \leq p - r \\
 c_i &= 2j + 3, & i &\in K_j, j \geq p - r + 1.
 \end{aligned}$$

INSERT FIGURE 10 ABOUT HERE

It is easy to verify that the algorithm outputs T_1 and f . Furthermore, since $c_i + c_j = 0$ if and only if $\{i,j\} = \{x,y\}$. The only other optimal solution is g by Remark 2.5. Therefore, f and g are adjacent extreme points of D_n . \square

We now develop a condition for adjacency of two threshold sequences that does not refer to their realizations as threshold graphs.

Lemma 3.6: Let $f = (f_1, \dots, f_n)$ be a threshold sequence with realization $T = (V, E_1)$ and $g = (g_1, \dots, g_n)$ be a degree sequence with realization $G = (V, E_2)$ such that $f = g + u_i + u_j$ for some $i \neq j$. (u_t is the t -th unit vector.) Then $[i,j] \in E_1$.

Proof: The proof is by induction on $\min(g_i, g_j)$. Let q be the number of positive components of g .

Basis: $\min(g_i, g_j) = 0$. Without loss of generality, $g_j = 0$. The number of positive components of f is $q + 1$ [$q + 2$] if g_i is positive [zero]. The largest degree in G is at most $q - 1$, and the largest degree in T is q [$q + 1$] by Theorem 1.3.

Since $\max_k g_k \leq q - 1$, we have $f_k \leq q - 1$ for $k \neq i, j$. Therefore, q [$q + 1$] = $\max(f_i, f_j)$, so one of the non-isolated vertices i, j of T has largest degree in T . Therefore, $[i, j] \in E_1$ by Theorem 1.3.

Induction step: $\min(g_i, g_j) > 0$. In this case the number of positive components of f is also q . If f_i or f_j is $q - 1$, then clearly $[i, j] \in E_1$. If not, $f_\ell = q - 1$ for some $\ell \neq i, j$ by Theorem 1.3. Then $g_\ell = q - 1$, so ℓ is adjacent to all non-isolated vertices in both T and G . Delete ℓ from

both T and G and apply induction to the resulting induced subgraphs. \square

Remark: One might wish to generalize Lemma 3.6 by relaxing one of its assumptions. Instead of assuming that $f = g + u_i + u_j$ for some $i \neq j$, we wish to assume only $f_k \geq g_k$ for all k , and to conclude that by dropping zero or more edges in the (unique) realization of f we can obtain some realization of g . Under the additional assumptions that (a) g is a threshold sequence and (b) $f_1 \geq \dots \geq f_n$ and $g_1 \geq \dots \geq g_n$, the conclusion is true [8]. However, (a) is not enough without (b), as can be seen from the example $f = (5,5,5,4,4,3)$, $g = (4,1,0,2,2,3)$.

Theorem 3.7: Let (f_1, \dots, f_n) and (g_1, \dots, g_n) be threshold sequences. Then they are adjacent extreme points of D_n if and only if there exist indices $i \neq j$ such that $f - g = \pm (u_i + u_j)$.

Proof: The "only if" part follows easily from Theorem 3.5.

("If): Assume without loss of generality that $f_k = g_k + 1$, $k = i, j$. From Lemma 3.6, $[i, j] \in E_1$, where $T_1 = (V, E_1)$ is the realization of (f_1, \dots, f_n) . From the uniqueness of realizations of threshold sequences, it follows that $T_2 = (V, E_1 - [i, j])$ is the realization of (g_1, g_2, \dots, g_n) , and the result follows from Theorem 3.5. \square

Below we obtain the threshold sequences adjacent to a given one without referring to its realization. For this it is convenient to use the following definition.

Definition 3.7 [7]: Let $f = (f_1, \dots, f_n)$ be any degree sequence. Let

$\delta_1 < \delta_2 \dots < \delta_m$ be the distinct nonzero integers occurring in f . Set $\delta_0 = 0$. The degree partition of f is defined as $D = (D(0), \dots, D(m))$ where

$$D(i) = \{j \mid f_j = \delta_i\}, \quad 0 \leq i \leq m.$$

we put

$$d_i = |D(i)|.$$

If f is a threshold sequence, then D is closely related to Definition 2.2. Indeed, if $G = (V, E)$ is the realization of f , let $P = (I_0, \dots, I_p, K_0, \dots, K_p)$ where K_i, I_i are as in Definition 2.2. A little reflection shows that if $|K_0| > 1$, then $D = P$, and if $|K_0| = 1$, then D is obtained from P by merging K_0 with I_p . Thus, if $|K_0| = 1$, then

$$D(i) = I_i, \quad 0 \leq i \leq p - 1$$

$$D(i) = K_0 \cup I_p, \quad i = p$$

$$D(i) = K_{i-p}, \quad p + 1 \leq i \leq m$$

$$m = 2p$$

and if $|K_0| > 1$, then

$$D(i) = I_i, \quad 0 \leq i \leq p$$

$$D(i) = K_{i-p-1}, \quad p + 1 \leq i \leq m$$

$$m = 2p + 1$$

Theorem 3.8: Let $f = (f_1, f_2, \dots, f_n)$ be a threshold sequence. Then the extreme points $f' = (f'_1, f'_2, \dots, f'_n)$ of D_n adjacent to f are given by:

$$(a) \quad f' = f + u_i + u_j$$

where $i \in D(r)$, $j \in D(m - r)$, $0 \leq r \leq \lfloor (m - 1)/2 \rfloor$

or $i, j \in D(\lfloor (m + 1)/2 \rfloor)$, $i \neq j$, m even;

$$(b) \quad f' = f - u_i - u_j$$

where $i \in D(r)$, $j \in D(m - r + 1)$, $1 \leq r \leq \lfloor m/2 \rfloor$

or $i, j \in D(\lfloor (m + 1)/2 \rfloor)$, $i \neq j$, m odd.

Proof: Let $T = (V, E)$ be the realization of f . Case (a) corresponds to adding an edge to E and Case (b) corresponds to dropping an edge from E . We prove the result for Case (a). Case (b) is similar.

First assume that m is odd. From Lemmas 3.2 and 3.3 we know that we can add an edge between I_r and K_{p-r} where $0 \leq r \leq p$ and no other edges. From Definition 3.7, this is the same as adding an edge between $D(r)$ and $D(m - r)$, $0 \leq r \leq \lfloor (m - 1)/2 \rfloor$.

Now assume that m is even. It follows from Lemmas 3.2 and 3.3 that we can add an edge between

- (1) I_r and K_{p-r} , $0 \leq r \leq p - 1$
- (2) I_p and K_0
- (3) any two vertices in I_p

As m is even, $|K_0| = 1$ and $K_0 \cup I_p = D(\lfloor (m + 1)/2 \rfloor)$. The result thus follows. \square

Using Theorem 3.8 we obtain the following result. We thank K. N. Srikanth for raising this question with one of us (M.S.).

Theorem 3.9: Let $f = (f_1, \dots, f_n)$ be a threshold sequence with a degree partition as in Definition 3.7. Then the number of extreme points of D_n adjacent to f is given by the expression

$$\sum_{i=0}^{\lfloor (m-1)/2 \rfloor} d_i d_{m-i} + \sum_{i=1}^{\lfloor m/2 \rfloor} d_i d_{m-i+1} + \binom{d_q}{2}$$

where $q = \lfloor (m-1)/2 \rfloor$. \square

We can now determine the threshold sequences with the largest and smallest number of adjacent sequences.

Theorem 3.10: Let $f = (f_1, f_2, \dots, f_n)$ be a threshold sequence with a degree partition as in Definition 3.7.

(a) The number of extreme points adjacent to f is a maximum, equal to $\binom{n}{2}$, if and only if

or $f_i = 0$, for all i (stable)

or $f_i = n - 1$, for all i (clique)

or $f_i = \begin{cases} n - 1 & \text{for some } j \\ 1 & \text{for } i \neq j \end{cases}$ (star)

or

$f_i = \begin{cases} 0 & \text{for some } i \\ n - 2 & \text{for } i \neq j \end{cases}$ (costar)

(b) The number of extreme points adjacent to f is a minimum, equal to n , if and only if

$$d_i = 1, i \neq 0, \lfloor (m + 1/2) \rfloor$$

$$d_i = 2, i = \lfloor (m + 1)/2 \rfloor$$

$$d_0 = n - \sum_{i=1}^m d_i.$$

Proof: Let $G = (V, E)$ be a realization of f . Let $V = K \cup I$ be as in Definition 2.2.

(a) The number of ways in which we can add or drop an edge preserving thresholdness is at most $\binom{n}{2}$. It is equal to $\binom{n}{2}$ if and only if every present edge can be dropped and every absent edge can be added. From Lemma 3.2 it follows that $p \leq 1$.

Case 1 ($p = 1$): It follows easily from Lemma 3.2 that in this case

$$|K_1| = 1, |K_0| = 1 \text{ and } I_0 = \emptyset. \text{ This implies that } G \text{ is a star.}$$

Case 2 ($p = 0$): We can similarly show that $\min(|K_0|, |I_0|) \leq 1$. This corresponds to G being a clique, a graph with no edges, or the complement of a star.

(b)

Case 1 (m is even): In this case $m = 2p$ and $d_p \geq 2$. From Theorem 3.9, the number of extreme points adjacent to f is

$$\begin{aligned} & \sum_{i=0}^{p-1} d_i d_{2p-i} + \sum_{i=1}^p d_i d_{2p-i+1} + \binom{d_p}{2} \\ &= d_0 d_{2p} + \sum_{i=1}^{p-1} d_i d_{2p-i} + \sum_{i=1}^p d_i d_{2p-i+1} + \binom{d_p}{2} \end{aligned}$$

$$\begin{aligned} &\geq d_0 + (p-1) + (p+1) + 1 \\ &= d_0 + 2p + 1 \end{aligned}$$

This lower bound is achieved if and only if

$$d_p = 2, d_i = 1, i \neq 0, p, d_0 = n - \sum_{i=1}^{2p} d_i.$$

The value of the bound is then $d_0 + 2p - 1 = (n - \sum_{i=1}^{2p} d_i) + 2p + 1$.

Case 2 (m is odd): In this case $m = 2p + 1$ and $d_{p+1} \geq 2$. the rest of the analysis is similar to Case 1. \square

4. Linear Description and Facets

In this section we use linear programming duality and Theorem 1.3 to provide a linear description of D_n .

Lemma 4.1: Let S, T be subsets of $\{1, 2, \dots, n\}$ such that $S \cap T = \emptyset$. Then the inequality

$$(4.1) \quad \sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$$

is valid for D_n . Moreover, the degree sequence of $G = (V, E)$ satisfies (4.1) with equality if and only if

- (a) $i, j \in S, i \neq j \Rightarrow [i, j] \in E$
- (b) $i, j \in T, i \neq j \Rightarrow [i, j] \notin E$

$$(c) \quad i \in S, j \in \overline{S \cup T} \Rightarrow [i, j] \in E$$

$$(d) \quad i \in T, j \in \overline{S \cup T} \Rightarrow [i, j] \notin E \quad (\text{See Figure 11}).$$

Proof: To show the validity of (1), we may assume that x is the degree sequence of $G = (V, E)$. Set $C = \{[i, j] | i \in S, j \in T\}$. It is easy to see that

$$(e) \quad \sum_{i \in S} x_i \leq |S| (n - 1 - |T| + |C|), \text{ equality holding if and only if} \\ (a) \text{ and } (c) \text{ hold, and}$$

$$(f) \quad -\sum_{i \in T} x_i \leq -|C|, \text{ equality holding if and only if } (b) \text{ and } (d) \text{ hold.}$$

By adding (e) and (f) we obtain (4.1) and the "moreover" part. \square .

INSERT FIGURE 11 ABOUT HERE

Theorem 4.2: D_n is determined by the linear inequalities

$$(4.2) \quad \sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$$

for all set S, T such that $\emptyset \neq S \cup T \subseteq \{1, 2, \dots, n\}$, $S \cap T = \emptyset$.

Proof: Let $c = (c_1, c_2, \dots, c_n) \in \mathbb{R}^n$, and consider the linear program

$$(4.3) \quad \max \sum_{i=1}^n c_i x_i \\ \text{subject to (4.2).}$$

Since it is known that Algorithm 2.3 maximizes $\sum_{i=1}^n c_i x_i$ over D_n , the theorem will be proved if it is shown that the solution produced by the algorithm on input c is an optimal solution of (4.3). Denote the graph produced by algorithm 2.3 by $G = (V, E)$ and its degree sequence by d . By Lemma 4.1, d is a feasible solution of (4.3).

The dual of (4.3) is:

$$(4.4) \quad \min \sum_{\substack{\emptyset \neq S \cup T \subseteq \{1, \dots, n\} \\ S \cap T = \emptyset}} |S| (n - 1 - |T|) y_{S, T}$$

s.t.

$$\sum_{\substack{\emptyset \neq S \cup T \subseteq \{1, \dots, n\} \\ S \cap T = \emptyset \\ i \in S}} y_{S, T} - \sum_{\substack{\emptyset \neq S \cup T \subseteq \{1, \dots, n\} \\ S \cap T = \emptyset \\ i \in T}} y_{S, T} = c_i, \quad i = 1, \dots, n$$

$$y_{S, T} \geq 0 \text{ for all } S, T \text{ such that } \emptyset \neq S \cup T \subseteq \{1, \dots, n\}, \\ S \cap T = \emptyset.$$

To show that d is an optimal solution of (4.3), it suffices to produce a feasible solution y to (4.4) that satisfies in addition the complementary slackness conditions:

$$(4.5) \quad y_{S, T} > 0 \text{ only if } \sum_{i \in S} d_i - \sum_{i \in T} d_i = |S| (n - 1 - |T|)$$

(this uses only weak duality; strong duality asserts that if in fact d is optimal to (4.3), then such a y exists; we do not need this result, as we construct y explicitly.)

Set $K = \{i | c_i \geq 0\}$ and $I = \{i | c_i < 0\}$, and put $r = |I|$. We may assume without loss of generality that $I = \{1, \dots, r\}$, $K = \{r + 1, \dots, n\}$,

$$c_{r+1} \leq c_{r+2} \leq \dots \leq c_n \text{ and } |c_r| \leq |c_{r-1}| \leq \dots \leq |c_1|$$

From the nature of Algorithm 2.3 it is clear that

$$(4.6) \quad N(1) \subseteq N(2) \subseteq \dots \subseteq N(r)$$

$$N(r+1) \cap I \subseteq N(r+2) \cap I \subseteq \dots \subseteq N(n) \cap I.$$

Consider the following pairs of sets (S_i, T_i) , $1 \leq i \leq n$:

$$(4.7) \quad S_i = \{i, i+1, \dots, n\} \quad r+1 \leq i \leq n$$

$$T_i = I - N(i)$$

and

$$S_i = N(i) \quad 1 \leq i \leq r$$

$$T_i = \{1, 2, \dots, i\}$$

Using Lemma 4.2 and (4.6), it can easily be verified that for the above choice of sets, S_i, T_i , $i = 1, \dots, n$, d satisfies the corresponding inequality in (4.2) with equality (see Figure 12). We will find a feasible solution y of (4.4) such that $y_{S,T} > 0$ only if $(S,T) = (S_i, T_i)$ for some i . The above two statements will then guarantee (4.5).

INSERT FIGURE 12 ABOUT HERE

Since $S_i \subseteq K$, $T_i \subseteq I$ for $1 \leq i \leq n$ and we imposed the condition $y_{S,T} = 0$ unless $(S,T) = (S_i, T_i)$ for some $1 \leq i \leq n$, the constraints (4.4) to be satisfied by $y_{S,T}$, $1 \leq i \leq n$, read:

$$(4.8) \quad \sum_{j \in T_i} y_{S_i, T_i} = |c_j|, \quad j = 1, \dots, r$$

$$\sum_{j \in S_i} y_{S_i, T_i} = c_j, \quad j = r + 1, \dots, n$$

$$y_{S_i, T_i} \geq 0, \quad i = 1, \dots, n.$$

For each $j \in \{1, \dots, n\}$, let P_j be the set of all indices i such that y_{S_i, T_i} appears in the j -th equation of (4.8). Then for $j \in I$ we have, by (4.6) and

(4.7)

$$(4.9) \quad \begin{aligned} P_j &= \{i \mid j \in T_i\} = \{i \in I \mid i \geq j\} \cup \{i \in K \mid [i, j] \notin E\} \\ &= \{i \in I \mid i \geq j\} \cup \{i \in K \mid c_i < |c_j|\}, \end{aligned}$$

and for $j \in K$, again by (4.6) and (4.7)

$$(4.10) \quad \begin{aligned} P_j &= \{i \in K \mid j \in S_i\} = \{i \in K \mid i \leq j\} \cup \{i \in I \mid [i, j] \in E\} \\ &= \{i \in K \mid i \leq j\} \cup \{i \in I \mid c_j \geq |c_i|\}. \end{aligned}$$

Define

$$\tilde{c}_t = \begin{cases} c_t & t \in K \\ |c_t| & t \in I \end{cases}$$

Let $\pi(1), \dots, \pi(n)$ be the rearrangement of $1, \dots, n$ such that

$$(a) \quad \tilde{c}_{\pi(1)} \leq \tilde{c}_{\pi(2)} \leq \dots \leq \tilde{c}_{\pi(n)};$$

(b) if $\tilde{c}_{\pi(i)} = \tilde{c}_{\pi(j)}$, $i \neq j$, then

$$(i - j)(\pi(i) - \pi(j)) \begin{cases} > 0 & \text{if } \pi(i), \pi(j) \in K \\ < 0 & \text{if } \pi(i), \pi(j) \in I \end{cases}$$

$\pi(i) < \pi(j)$ if $\pi(i) \in I$, $\pi(j) \in K$.

Then $P_{\pi(j)} = \{\pi(1), \dots, \pi(j)\}$ by (4.9) and (4.10).

We can now rearrange the n equations (4.8) so that the t -th new equation is the $\pi(t)$ -th equation of (4.8), and also arrange the n variables $y_{S_1, T_1}, \dots, y_{S_n, T_n}$ so that the t -th new variable is $y_{S_{\pi(t)}, T_{\pi(t)}}$. Then the new system has the form $Ay = b$ where the coefficient matrix A has zeros above the main diagonal and ones everywhere else, and the components of b are nonnegative and nonincreasing. This system has a unique, nonnegative solution y . This completes the proof. \square

Corollary 4.3: The system (4.2) of linear inequalities is totally dual integral; that is, whenever c_1, \dots, c_n are integers, the dual of (4.3) has an optimum solution in integers. \square

For more information on totally dual integral systems, see [4, 20, 21].

In order to identify the facets of D_n , we need to know its dimension and that of a related polytope. For $m \geq 1$, $n \geq 1$, let $K_{m,n}$ be the complete bipartite graph with dipartition $\{1, 2, \dots, m\}$, $\{m+1, m+2, \dots, m+n\}$, and let $D_{m,n}$ be the convex hull of the degree sequences of the spanning subgraphs of $K_{m,n}$ (just as D_n is the convex hull of the degree sequences of

the spanning subgraphs of K_n , the complete graph on $\{1, \dots, n\}$.

Lemma 4.3:

- (a) $\dim D_1 = 0$, $\dim D_2 = 1$;
- (b) $\dim D_{m,n} = m + n - 1$ for $m \geq 1$, $n \geq 1$;
- (c) $\dim D_n = n$ for $n \geq 3$.

Proof:

- (a) Clear.
- (b) Since $D_{m,n}$ lies on the hyperplane $\sum_{i=1}^m x_i = \sum_{i=m+1}^{m+n} x_i$, we have $\dim D_{m,n} \leq m + n - 1$. To prove the opposite inequality, consider any spanning tree of $K_{m,n}$ and let A be its edge-vertex incidence matrix. The $m + n - 1$ rows of A are linearly independent, and each of them lies in $D_{m,n}$. Hence $\dim D_{m,n} \geq m + n - 1$.
- (c) Koren [15] proved (c) by showing that D_n has interior points for $n \geq 3$. We prove it as follows: for n odd, let H be any Hamiltonian cycle in K_n , and for n even, let H consist of any cycle through $n - 1$ vertices and a single edge at the remaining vertex. Let A be the edge-vertex incidence matrix of H . In each case A is nonsingular and its rows lie in D_n . Hence $\dim D_n \geq n$. \square

Theorem 4.4:

For $n \geq 3$, the facets of D_n are given by:

- (a) $x_i \geq 0$, $i = 1, \dots, n$ (only if $n \geq 4$).
- (b) $x_i \leq n - 1$, $i = 1, \dots, n$ (only if $n \geq 4$).

- (c) $\sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$ for all sets S, T such that
 $S \neq \emptyset, T \neq \emptyset, S \cap T = \emptyset, S \cup T \subseteq \{1, \dots, n\}, |S \cup T|$
 $= 2, 3, \dots, n - 3, n.$

Proof:

By Theorem 4.2, every facet of D_n has the form

$$(4.11) \quad \sum_{i \in S} x_i - \sum_{i \in T} x_i \leq |S| (n - 1 - |T|)$$

for some sets S, T with $\emptyset \neq S \cup T \subseteq \{1, \dots, n\}, S \cap T = \emptyset.$

Let $D_{S,T}$ denote the set of degree sequences satisfying (4.11) with equality. Thus by (c) of Lemma 4.3, (4.11) is a facet of D_n if and only if $\dim D_{S,T} = n - 1.$ From Lemma 4.1 it follows that $\dim D_{S,T} = \dim D_{|S|, |T|} + \dim D_{|\overline{S \cup T}|},$ where $\dim D_{|S|, |T|}$ means 0 if either S or T vanishes. Thus for $S = \emptyset,$ (4.11) is a facet if and only if $|\bar{T}| = n - 1$ and $|\bar{T}| \geq 3,$ which corresponds to (a). Similarly for $T = \emptyset,$ (4.11) is a facet if and only if $|\bar{S}| = n - 1$ and $|\bar{S}| \geq 3,$ which corresponds to (b). Finally for $S \neq \emptyset, T \neq \emptyset,$ (4.11) is a facet if and only if $|S| + |T| - 1 + \dim D_{|\overline{S \cup T}|} = n - 1,$ or equivalently $\dim D_{|\overline{S \cup T}|} = |\overline{S \cup T}|,$ which holds if and only if $|\overline{S \cup T}| = 0, 3, 4, \dots, n$ by Lemma 4.3. This case corresponds to (c). \square

5. Threshold Sequences and Majorization

Definition 5.1: For any real sequence $f = (f_1, \dots, f_n),$ let

$f_{[1]} \geq f_{[2]} \geq \dots \geq f_{[n]}$ denote the components of f sorted in a

nonincreasing order. We say that f majorizes $g,$ denoted $f \succeq g,$ when

$\sum_{i=1}^k f_{[i]} \geq \sum_{i=1}^k g_{[i]}$ holds for each $k = 1, \dots, n$, with equality holding for $k = n$. We say that f strictly majorizes g , denoted $f > g$, when $f \succeq g$ and at least one of the above inequalities for $k = 1, \dots, n - 1$ is strict. This means that the common sum of the components of f and g is distributed more evenly in g than in f .

Remark 5.2:

- (a) If f and g majorize each other, then f is a rearrangement of g , and conversely.
- (b) If $f > g$, then f is not a rearrangement of g .
- (c) Majorization and strict majorization are transitive.
- (d) For a wealth of characterizations and applications of majorization, see [12, 17]. [17] denotes majorization by $>$, not by \succeq .

Definition 5.2: Let $f = (f_1, f_2, \dots, f_n)$ be an integer sequence and assume that $f_i \geq f_j + 2$ for some i, j . Then the sequence $g = f - u_i + u_j$ is said to be obtained from f by a unit transformation from i to j (u_t is the t -th unit vector). In that case, clearly $f > g$. Conversely,

Theorem 5.3 [12, 17]: If f and g are integer sequences such that $f \succeq g$, then some rearrangement of g can be obtained from f by a sequence (possibly empty) of unit transformations. \square

Definition 5.4: Let $G = (V, E)$ be a graph and let $i, j, k \in V$ be such that $\deg k \geq \deg i + 2$, $[i, j] \notin E$, $[j, k] \in E$. Then the operation of dropping

from G the edge $[j,k]$ and adding the edge $[i,j]$ is called a just rotation from k to i (see Figure 13).

INSERT FIGURE 13 ABOUT HERE

The adjective "just" is used in the sense of the "rich" vertex (k) giving some of its degree to the "poor" one (i). Thus, the degree sequence of G strictly majorizes the resulting degree sequence.

Lemma 5.5: A just rotation cannot result in a threshold graph.

Proof: Note that $\deg k \geq \deg i$ in the resulting graph, yet i has a neighbor j that is not a neighbor of k . This would contradict Theorem 1.3 if the resulting graph were threshold. \square

Lemma 5.6: Let d be the degree sequence of a graph $G = (V,E)$, and let d' result from d by a unit transformation. Then some just rotation in G results in a graph with degree sequence d' .

Proof: Let d' result from d by a unit transformation from k to i . Then $d'_k \geq d_i + 2$ in G , and therefore there is a vertex j such that $[i,j] \notin E$, $[j,k] \in E$. Hence G has a just rotation from k to i , resulting in a graph with degree sequence d' . \square

We now characterize the threshold sequences by the property of non-majorizability.

Theorem 5.7: A degree sequence d is threshold if and only if d is not strictly majorized by any degree sequence.

Proof: ("If"): Assume d is the degree sequence of the non-threshold graph $G = (V, E)$. By Corollary 1.2, G has vertices i, j, k, ℓ with $[i, j], [\ell, k] \in E$, $[i, \ell], [k, j] \notin E$ (Figure 14).

INSERT FIGURE 14 ABOUT HERE

Without loss of generality, $\deg i \leq \deg k$ in G . Let the graph H result from G by deleting edge $[i, j]$ and adding edge $[j, k]$. The reverse operation is then a just rotation in H , hence the degree sequence of H strictly majorizes d .

("Only if"): Assume $f > d$ and let G realize f . By Theorem 5.3, there are sequences $f^{(0)}, f^{(1)}, \dots, f^{(t)}$ such that $f^{(0)} = f$, $f^{(t)}$ is a rearrangement of d , and some unit transformation on $f^{(i)}$ produces $f^{(i+1)}$ for $i = 0, \dots, t - 1$. By Lemma 5.6 it follows that some sequence of just rotations in G results in a graph with degree sequence $f^{(t)}$. By Lemma 5.3, $f^{(t)}$ is a non-threshold sequence. Therefore, d is a non-threshold sequence. \square

Corollary 5.8: Every degree sequence is a convex combination of the rearrangements of a single threshold sequence (i.e., of the degree sequences of isomorphic threshold graphs).

Proof: Let d be a threshold sequence. If d is a threshold, there is nothing to prove. Otherwise, d is strictly majorized by some degree sequence d' by Theorem 5.7. Continue this process as long as possible. The progression d, d', d'', \dots has no repetitions by Remark 5.2 because each term is strictly majorized by the next one. It must terminate, because each of its terms is a sequence of n nonnegative integers with the same sum. By Theorem 5.7, the last sequence f is threshold and it majorizes d by transitivity. The result then follows from the well-known fact that f majorizes d if and only if d is a convex combination of the rearrangements of f [12, 17]. \square

References

- [1] J. S. Beissinger and U. N. Peled, "Enumeration of Labelled Threshold Graphs and a Theorem of Frobenius Involving Eulerian Polynomials," Graphs and Combinatorics, 3 (1987), 213-219.
- [2] C. Berge, Graphs and Hypergraphs, Amsterdam: North Holland, 1973.
- [3] V. Chvátal and P. L. Hammer, "Aggregation of Inequalities in Integer Programming," Ann. Discr. Math., 1 (1977), 145-162.
- [4] J. Edmonds and R. Giles, "Total Dual Integrality of Linear Inequality Systems," in Progress in Combinatorial Optimization, W. R. Pulleyblank (ed.), Academic Press, 1984.
- [5] P. Erdős and T. Gallai, "Graphs with Prescribed Degrees of Vertices" (Hungarian), Mat. Lapok II (1960), 264-272.
- [6] M. C. Golumbic, "Threshold Graphs and Synchronizing Parallel Processes," Colloq. Math. Soc. J. Bolyai (Combinatorics), 18 (1978), 419-428.
- [7] M. C. Golumbic, Algorithmic Graph Theory and Perfect Graphs, Academic Press, 1980.
- [8] S. L. Hakimi, "On the Realizability of a Set of Integers as Degrees of the Vertices of a Graph," SIAM J. Appl. Math., 10 (1960), 496-506.
- [9] P. L. Hammer, T. Ibaraki and B. Simeone, "Threshold Sequences," SIAM J. Alg. Discr. Meth., 2 (1981), 39-49.
- [10] P. L. Hammer and B. Simeone, "The Splittance of a Graph," Combinatorica, 1 (1981), 275-284.
- [11] F. Harary and U. N. Peled, "Hamiltonian Threshold Graphs," Disc. Appl. Math., 16 (1987), 11-15.

- [12] G. H. Hardy, J. E. Littlewood and G. Pólya, Inequalities, Cambridge University Press, 1959.
- [13] V. Havel, "A Remark on the Existence of Finite Graphs" (Czech), Časopis Pěst. Mat., 80 (1955), 477-480.
- [14] P. B. Henderson and Y. Zalcstein, "A Graph-theoretic Characterization of the PV Chunk Class of Synchronizing Primitives," SIAM J. Comput., 6 (1977), 88-108.
- [15] M. Koren, "Extreme Degree Sequences of Simple Graphs," J. Comb. Theory (B), 15 (1973), 213-224.
- [16] L. Lovász, Combinatorial Problems and Exercises, North Holland, 1979.
- [17] A. W. Marshall and I. Olkin, Inequalities: Theory of Majorization and its Applications, Academic Press, 1979.
- [18] J. Orlin, "The Minimal Integral Separator of a Threshold Graph," Ann. Discr. Math., 1 (1977), 415-419.
- [19] U. N. Peled, "Threshold Graph Enumeration and Series-product Identities," Congressus Numeratum, 29 (1980), 735-738.
- [20] W. R. Pulleyblank, "Polyhedral Combinatorics," in Mathematical Programming--the State of the Art, Bonn 1982, A. Bachem, M. Grötschel and B. Korte (eds.), Springer-Verlag, 1983.
- [21] A. Schrijver, Theory of Linear and Integer Programming, John Wiley and Sons, 1986.
- [22] D. J. Welsh, Matroid Theory, Academic Press, 1976.




Fig. 1: The forbidden induced subgraphs of threshold graphs.




Fig. 2: The forbidden configuration of threshold graphs.




Fig. 3: A typical threshold graph. A line between K_i and I_j (or between K_i and K_j) indicates that every vertex of K_i is adjacent to every vertex of I_j (K_j).




Fig. 4: Illustrating the objective function for the "if" part of Theorem 2.8.




Fig. 5: An interchange.




Fig. 6: Illustrating the proof of Lemma 3.2.




Fig. 7: Illustrating the proof of Lemma 3.2.




Fig. 8: Illustrating the proof of Lemma 3.2.




Fig. 9: Illustrating the proof of Lemma 3.3.




Fig. 10: Weights in the proof of Theorem 3.5.




Fig. 11: S: Clique. All edges between S and \overline{SUT} present;
T: Stable. All edges between T and \overline{SUT} absent.




Fig. 12: Illustrating the choice of dual variables in the proof of Theorem 4.2.




Fig. 13: A just rotation.




Fig. 14: Illustrating the "if" part of Theorem 5.6.