

# Common Belief Foundations of Global Games\*

Stephen Morris  
Princeton University

Hyun Song Shin  
Bank for International Settlements

Muhamet Yildiz  
M.I.T.

April 2015

## Abstract

This paper makes two related contributions to the study of Global Games. First, for general type spaces, it characterizes rationalizable actions in terms of the properties of the belief hierarchies and shows that there is a unique rationalizable action played whenever there is approximate common certainty of rank beliefs, defined as the probability the players assign to their payoff parameters being higher than their opponents'. We argue that this is the driving force behind selection results for the specific type spaces in the global games literature. Second, applications of global games often assume players' payoff parameters are derived from common and idiosyncratic terms with normal distributions. This paper analyzes the alternative case that these terms have fat-tailed distributions. If the common component has a thicker tails, then there is common certainty of rank beliefs, and thus uniqueness and equilibrium selection. If the idiosyncratic terms has thicker tails, then there is approximate common certainty of payoffs, and thus multiple rationalizable actions.

## 1 Introduction

Complete information games often have many equilibria. Even when they have a single equilibrium, they often have many actions that are rationalizable, and are therefore consistent with common certainty of rationality. The inability of theory to make a prediction is problematic for economic applications of game theory.

Carlsson and van Damme (1993) suggested a natural perturbation of complete information that gives rise to a unique rationalizable action for each player. They introduced the idea of “global games”—where any payoffs of the game are possible and each player observes the true payoffs of the game with a small amount of noise. They showed—for the case of two player two action games—that as the noise about payoffs

---

\*This paper incorporates material from a working paper of the same title circulated in 2007 (Morris and Shin (2007)). We are grateful for comments from seminar participants at Columbia and Northwestern on this iteration of the project. The views expressed in this paper reflects those of the authors and not necessarily those of the Bank for International Settlements.

become small, there is a unique equilibrium; moreover, the perturbation selects a particular equilibrium (the *risk dominant* one) of the underlying game. This result has since been generalized in a number of directions and widely used in applications.<sup>1</sup> When the global game approach can be applied to more general games, it can be used to derive unique predictions in settings where the underlying complete information game has multiple equilibria, making it possible to carry out comparative static and policy analysis. An informal argument that has been made for this approach is that the "complete information" assumption is an unrealistic one that makes it easier to support multiplicity, so the natural perturbation underlying global games captures the more realistic case.

However, the global game selection result uses a particular form of perturbation away from "complete information." Complete information entails the assumption that a player is certain of the payoffs of the game, certain that other players are certain, and so on. Weinstein and Yildiz (2007) consider more general perturbations, saying that a situation is close to a complete information game if players are almost certain that payoffs are close to those complete information game payoffs, almost certain that other players are almost certain that payoffs are close to those payoffs, and so on. Formally, they consider closeness in the product topology on the universal belief space. They show that for *any* action which is rationalizable for a player in a complete information game, there exists a nearby type of that player in the product topology for whom this is the unique rationalizable action. Thus by considering a richer but also intuitive class of perturbations, they replicate the global game uniqueness result but reverse the selection result.

In this paper, we identify the driving force behind global game uniqueness and selection results. In particular, we do not want to take literally the (implicit) assumption in global games that there is common certainty among the players of a technology which generates (conditionally independent) noisy signals observed by the players. Rather, we want to argue that global game perturbations are a metaphor, or a convenient modelling device, for a more general intuitive class of relaxations of common certainty. We want to characterize and analyze the key property of that more general class, which must also be more restriction than the product topology perturbations of Weinstein and Yildiz (2007).

Our analysis is carried out for a two player, binary action game. Each player must decide whether to "invest" or "not invest". Payoffs are given by the following matrix:

	invest	not invest	
invest	$x_1, x_2$	$x_1 - 1, 0$	(1)
not invest	$0, x_2 - 1$	$0, 0$	

Each player  $i$  knows his own payoff parameter, or return to investment,  $x_i$  but may not know the other

---

<sup>1</sup>Morris and Shin (1998) analyzed a global game with a continuum of players making binary choices, and this case has been studied in a number of later applications. See Morris and Shin (2003b) for an early survey of some theory and applications of global games. Frankel, Morris, and Pauzner (2003) study global game selection in general games with strategic complementarities.

player's payoff parameter. There are strategic complementarities, because a player has a loss of 1 if the other player does not invest. If  $x_1$  and  $x_2$  are both in the interval  $[0, 1]$ , then there are multiple equilibria, both invest and both not invest, under complete information. In the symmetric case, with  $x_1 = x_2 = x$ , the *risk dominant* equilibrium in this game is the equilibrium that has each player choose a best response to a 50/50 distribution over his opponent's action. Thus both invest is the risk dominant equilibrium if  $x > \frac{1}{2}$ .

If  $x_1$  and  $x_2$  are both in the interval  $[0, 1]$ , both actions remain rationalizable for player  $i$  if there is approximate common certainty of payoffs. There is approximate common certainty of an event if both believe it with probability close to 1, both believe with probability close to 1 that both believe it with probability close to 1, and so on. This is a well known sufficient condition for multiple rationalizable actions, going back to Monderer and Samet (1989). But it is a strong condition. A key concept to understand uniqueness in this setting is a player's "rank belief"—that is, his belief about whether he has the higher payoff parameter (so he has rank 1) or the lower payoff parameter (so he has rank 2). A player has the *uniform rank belief* if he assigns probability  $\frac{1}{2}$  to each rank. If there is *common certainty of uniform rank beliefs*, then a player has a unique rationalizable action. In particular, action invest is uniquely rationalizable for a player if it is risk dominant. To see why this is true, let  $x^*$  be the smallest payoff parameter such that invest is uniquely rationalizable whenever there is common certainty of uniform rank beliefs for a player with payoff parameter greater than or equal to  $x^*$ . If  $x^*$  were strictly greater than  $\frac{1}{2}$ , a player with payoff parameter close to  $x^*$  (and common certainty of uniform rank beliefs) would assign probability close to  $\frac{1}{2}$  to his opponent investing and would therefore have a strict incentive to invest. Thus we would have a contradiction. Thus  $x^*$  must be less than or equal to  $\frac{1}{2}$ . This proves the result. This is the standard global game argument. But because it is expressed for general situations—and, in particular, without reference to noisy signals or one dimensional type spaces—it provides a primitive common belief foundation for global game selection. One of our main results will be a formalization of an appropriate weakening of this sufficient condition: *approximate* common certainty of *approximately* uniform rank beliefs.

A second contribution of this paper is to identify new and insightful conditions under which either approximate common certainty of payoffs and or approximate common certainty of approximately uniform rank beliefs will emerge. Like the classical global games literature, we will consider the case where players have one dimensional types that reflect both common and idiosyncratic components, and consider sequences approaching complete information. Unlike the classical global games literature, we consider the case where the common and idiosyncratic components have distributions with "fat tails," i.e., with greater than exponential densities. Fat tails generate intuitive and insightful higher-order beliefs as complete information is approached. In particular, close to the complete information limit, if a player observes a payoff parameter which is surprisingly different from his prior expectation, does he attribute the difference to the common component or the idiosyncratic component? Under fat tails, the most likely explanation

is that either one or the other but not both had an extreme realization. If the idiosyncratic component has a fatter tail, then a player attributes the difference to the idiosyncratic component. He thinks that the common component was close to its mean, and thus expects that his opponent's payoff type is closer to that mean. Thus, as we approach the complete information limit, there is approximate common certainty of payoffs. In that case, there will be multiple rationalizable actions whenever there are multiple equilibria under complete information. But if the common component has a fatter tail, then he thinks that an unusual realization of the common component accounts for the deviation, and thus he expects his opponent's payoff type to be close to his. He finds it equally likely that either player has the higher signal. Thus there is approximate common certainty of uniform rank beliefs. This leads to unique selection of risk-dominant equilibrium as in the global games literature. Thus global game uniqueness and selection is tied to whether the players' idiosyncratic component has a thinner tail.

We can contrast these results to existing results in the global games literature. The leading exercise in the global games literature is to consider a fixed prior about payoffs and have players observe conditionally independent noise about payoffs (Carlsson and van Damme (1993), Morris and Shin (2003b) and Frankel, Morris, and Pauzner (2003)). As the noise shrinks to zero, there is a unique equilibrium. The arguments for these results rely on uniform convergence of conditional probability properties which, in the case of symmetric settings, reduces to (variants of) common certainty of rank beliefs. Another leading exercise is to examine what happens if there is normally distributed common payoff parameter, and players observe the prior mean (a public signal) and also a conditionally independent normal private signal (Morris and Shin (2001) and Morris and Shin (2003a)), and allow the variances of both the prior and the noisy private signals to converge to zero. In this case, there is a unique equilibrium if and only if the noise in the private signal shrinks sufficiently fast relative to the public signal. Intuitively, relatively fast convergence of private noise gives common certainty of rank beliefs in the limit while relatively slow convergence gives approximate common certainty of payoffs.<sup>2</sup> The critical condition turns out to be that the variance of the private signal must shrink faster than the square of the variance of the public signal. However, this rate condition is special to the normal distribution and our results highlight quite different properties that determine limit behavior in general. We find the fat tail result more insightful and robust than the normal distribution insights which two of us have highlighted in earlier work. This paper is the first to examine what happens to higher-order beliefs or global game results without normality when both the prior and private signals become concentrated.

These results shed light on when global game models can offer useful insights in applications. The one dimensional normal model suggested the simple message that there will be unique rationalizable outcomes if

---

<sup>2</sup>As we will show in the body of the paper, common certainty of rank beliefs in the limit never arises in the normal case, but a weaker condition - common certainty of rank beliefs conditional on undominated actions - is satisfied and sufficient for limit uniqueness.

there is enough private information (measured by its precision) relative to the amount of public information. This suggests that we can assess the relevance of the uniqueness prediction by focussing on whether private signals have less variance than public signals (see, e.g., Svensson (2006)); and it suggests that uniqueness results will be less relevant in settings where information is being endogenously generated in a public way (see, e.g., Atkeson (2001)).<sup>3</sup> Going beyond the one dimensional normal model (where precision is the unique parameter characterizing the shape of the distribution) suggests more nuanced conclusions. Our fat tail results highlight the importance of the shape of the tail of the distribution for determining the relevant properties of higher-order beliefs. And, more generally, the common belief foundations results focus attention on the properties of higher-order beliefs that matter for global game results rather than the conditionally independent noisy signal story that generates them.

We present our results in the context of the simple, two player, strategic setting described above, maintaining the common prior assumption. We do this in order to focus on the common belief foundations rather than the details of the strategic setting. But we can then describe how our results can be mapped back to more general settings considered in the global games literature.

Under the common prior assumption, if there is common certainty that players' beliefs about their rank is  $r \in [0, 1]$ , we must have  $r = \frac{1}{2}$ , i.e., what we refer to as uniform rank beliefs. However, if the common prior assumption is dropped, common certainty of rank beliefs continues to lead to uniqueness and selection, but the selected action now depends on the rank belief  $r$ . This observation provides common belief foundations for the results of Izmalkov and Yildiz (2010).

We state our results for symmetric two player two action games, but they extend to symmetric many player two action games. The common belief characterization of rationalizability becomes more complex, because the relevant belief operators depend on beliefs about more than one event. However, the limit uniqueness and multiplicity results extend cleanly. Morris and Shin (2003b) highlight that in many player symmetric games, the selected equilibrium is the *Laplacian* one: each player is choosing a best response to a uniform conjecture over the number of his opponents choosing each action. This selection is key in the vast majority of applied papers. Common certainty of rank beliefs in the  $N$  player case corresponds to assigning probability  $\frac{1}{N}$  to exactly  $n$  players having higher payoff states, for each  $n = 0, 1, \dots, N - 1$ , and is a sufficient condition for the global game selection of the Laplacian action. Multiplicity when there is approximate common certainty of payoffs holds for general games, as shown by Monderer and Samet (1989).

Most of the global games literature focusses on a "common values" case, where there is a common payoff parameter of interest and each player observes it with idiosyncratic private noise. Our analysis concerns a "private value" case, where the idiosyncratic shock is payoff relevant to the players. Results in

---

<sup>3</sup>See also Angeletos, Hellwig, and Pavan (2007), Angeletos and Werning (2006), Hellwig (2002) and Hellwig, Mukerji, and Tsyvinski (2006).

the private value case are close to those in the common value case when noise is small (see Morris and Shin (2005) for some theory and Argenziano (2008) for an application of private value global games). Analogous results could be obtained for the common value case, but at the expense of significantly more complicated statements to account for the small amount of payoff uncertainty. Another convenient property of the game we consider is the additive separability between a component of payoffs that depends on others' actions and a component about which there is uncertainty. This gives a sharp separation between the role of strategic uncertainty and payoff uncertainty. Again, this provides sharper results although qualitative similar results would go through without this assumption.

However, the global game selection results reported in the paper rely on symmetric payoffs; in particular, it is required for common certainty of approximately uniform rank beliefs to be the relevant sufficient condition for unique rationalizable actions. Higher-order belief foundations can be provided for asymmetric global game results, but they are qualitatively different from those provided for symmetric games here.

We present a general characterization of rationalizable behavior in Section 2, reporting sufficient conditions, in general type spaces, for multiple rationalizable outcomes based on approximate common certainty of payoffs and for unique rationalizable outcomes based on approximate common certainty of approximately uniform rank beliefs. In Section 3, we introduce and analyze the one dimensional model generated by common and idiosyncratic payoff shocks with fat tails. In Section 4, we discuss extensions to more general strategic settings and the relation to the global games literature.

## 2 General Type Spaces

### 2.1 Higher Order Beliefs

There are two players, 1 and 2. Let  $T_1$  and  $T_2$  be the sets of types for players 1 and 2, respectively. A mapping  $x_i : T_i \rightarrow \mathbb{R}$  describes a payoff parameter of interest to player  $i$  and a mapping  $\pi_i : T_i \rightarrow \Delta(T_j)$  describes player  $i$ 's beliefs about the other player. We assume that  $T_i$  is endowed with a topology and Borel sigma-algebra. We make a couple of minimal continuity assumptions:  $x_i$  and  $\pi_i$  are continuous, and the pre-image  $x_i^{-1}([a, b])$  of every compact interval  $[a, b]$  is sequentially compact. This type space can be arbitrarily rich, and in particular can encode any beliefs and higher-order beliefs, and thus our results apply if the type space is the universal (private value) belief space of Mertens and Zamir (1985).<sup>4</sup>

We start with describing the belief and common belief operators, as in Monderer and Samet (1989).

---

<sup>4</sup>Mertens and Zamir (1985) constructed a space that encodes all beliefs and higher order beliefs about a common state space. We maintain the assumption that the state space is a pair of payoff types of the players, where each player knows his own payoff type. It is a simple adaption to the classic construction to build in this restriction, see for example Heifetz and Neeman (2006). The classical construction assumes a compact state space. We need to allow the state space to be  $\mathbb{R}^2$  but instead impose sequential compactness.

The state space is  $T = T_1 \times T_2$ . An *event* is a subset of  $T$ . An event is *simple* if  $E = E_1 \times E_2$  where  $E_i \subseteq T_i$ . For our game theoretic analysis, we will be interested in events that are simple and compact and we restrict attention to such events in the analysis that follows. For any such simple and compact event  $E$ , we write  $E_1$  and  $E_2$  for the compact projections of  $E$  onto  $T_1$  and  $T_2$ , respectively. Now, for probability  $p_i$ , write  $B_i^{p_i}(E)$  for the set of states where player  $i$  believes  $E$  with probability at least  $p_i$ :

$$B_i^{p_i}(E) = \{(t_1, t_2) \mid t_i \in E_i \text{ and } \pi_i(E_j \mid t_i) \geq p_i\}.$$

For a pair of probabilities  $(p_1, p_2)$ , say that event  $E$  is  $(p_1, p_2)$ -*believed* if each player  $i$  believes event  $E$  with probability at least  $p_i$ . Writing  $B_*^{p_1, p_2}(E)$  for the set of states where  $E$  is  $(p_1, p_2)$ -believed, we have:

$$B_*^{p_1, p_2}(E) = B_1^{p_1}(E) \cap B_2^{p_2}(E).$$

Say that there is *common*  $(p_1, p_2)$ -*belief* of event  $E$  if it is  $(p_1, p_2)$ -believed, it is  $(p_1, p_2)$ -believed that it is  $(p_1, p_2)$ -believed, and so on. We write  $C^{p_1, p_2}(E)$  for set of states at which  $E$  is common  $(p_1, p_2)$ -belief. Thus

$$C^{p_1, p_2}(E) = \bigcap_{n \geq 1} [B_*^{p_1, p_2}]^n(E).$$

An event is  $(p_1, p_2)$ -*evident* if it is  $(p_1, p_2)$ -believed whenever it is true. Generalizing a characterization of common knowledge by Aumann (1976), Monderer and Samet (1989) provides the following useful characterization.

**Lemma 1 (Monderer and Samet (1989))** *Event  $E$  is common  $(p_1, p_2)$ -belief if and only if there exists a  $(p_1, p_2)$ -evident event  $F$  with  $F \subseteq B_*^{p_1, p_2}(E)$ .*

In our formulation, we make the belief operators type dependent, and the above properties generalize immediately to this case. For any  $f_i : T_i \rightarrow \mathbb{R}$ , we say that event  $E$  is  $f_i$ -*believed by type  $t_i$*  of player  $i$  if he believes it with probability at least  $f_i(t_i)$ :

$$B_i^{f_i}(E) = \{(t_1, t_2) \mid t_i \in E_i \text{ and } \pi_i(E_j \mid t_i) \geq f_i(t_i)\}.$$

Clearly we can make richer statements about beliefs and higher-order beliefs in this language. We will continue to write  $B_i^{p_i}(E)$  for the original  $p_i$ -belief operator, where  $p_i$  is now understood as the constant function of types taking the value  $p_i$ . Note that we allow  $f_i$  to take values below 0 and above 1. This convention gives a special role to the events  $B_i^{f_i}(\emptyset)$  and  $B_i^{f_i}(T)$ , since a player always believes an event with probability at least 0 and never believes an event with probability greater than 1. Thus

$$\begin{aligned} B_i^{f_i}(\emptyset) &= \{(t_1, t_2) \mid f_i(t_i) \leq 0\} \\ B_i^{f_i}(T) &= \{(t_1, t_2) \mid f_i(t_i) \leq 1\}. \end{aligned}$$

These operators behave just like the type-independent ones. In particular, writing  $B_*^{f_1, f_2}(E)$  for the set of states where  $E$  is  $(f_1, f_2)$ -believed, we have:

$$B_*^{f_1, f_2}(E) = B_1^{f_1}(E) \cap B_2^{f_2}(E).$$

Say that there is *common*  $(f_1, f_2)$ -belief of event  $E$  if it is  $(f_1, f_2)$ -believed, it is  $(f_1, f_2)$ -believed that it is  $(f_1, f_2)$ -believed, and so on. We write  $C^{f_1, f_2}(E)$  for set of states  $E$  is common  $(f_1, f_2)$ -belief. An event is  $(f_1, f_2)$ -evident if it is  $(f_1, f_2)$ -believed whenever it is true:

$$C^{f_1, f_2}(E) = \bigcap_{n \geq 1} \left[ B_*^{f_1, f_2} \right]^n (E)$$

The Monderer and Samet (1989) characterization immediately generalizes to our case.

**Lemma 2** *Event  $E$  is common  $(f_1, f_2)$ -belief if and only if there exists a  $(f_1, f_2)$ -evident event  $F$  with  $F \subseteq B_*^{f_1, f_2}(E)$ .*

## 2.2 Common-Belief Characterization of Rationalizability

In the baseline model, we consider the following action space and the payoff function:

	invest	not invest
invest	$x_1, x_2$	$x_1 - 1, 0$
not invest	$0, x_2 - 1$	$0, 0$

Each  $x_i$  is assumed to be a continuous function of type  $t_i$ . Player  $i$  knows his own payoff parameter  $x_i(t_i)$  but does not necessarily know the other player's payoff parameter  $x_j(t_j)$ . Moreover, he gets a return  $x_i(t_i)$  if he invests but faces a penalty 1 if the other player does not invest. Hence, he only wants to invest if the probability he assigns to his opponent investing is at least  $1 - x_i(t_i)$ . We now define rationalizability in the context of this game (it corresponds to standard general definitions). Say that an action is  $(k + 1)$ th level rationalizable if it is a best response to  $k$ th level rationalizable play of his opponent; and say that any action is 0th level rationalizable. Write  $R_i^k$  for the set of types of player  $i$  for whom action invest is  $k$ th-level rationalizable and let  $R_i^0 = T_i$ .

We start with carefully describing the set  $R_i^1$  of types for whom invest is 1st-level rationalizable in terms of our type-dependent belief operators. On the one hand, for any type  $t_i$  of player  $i$ , action invest is 1st-level rationalizable for  $t_i$  if and only if  $x_i(t_i) \geq 0$ ; this is in response to the belief that his opponent invests with probability 1. On the other hand, since player  $i$  always assigns probability 1 to  $T$ , he assigns probability at least  $1 - x_i(t_i)$  to  $T$  if and only if  $x_i(t_i) \geq 0$ . Thus,

$$R_i^1 = B_i^{1-x_i}(T).$$



That is, the types for which invest is first-level rationalizable coincide with those in  $B_i^{1-x_i}(T)$ .

Now, action invest is 2nd-level rationalizable for type  $t_i$  of player  $i$  if, in addition, he assigns probability at least  $1 - x_i(t_i)$  to  $x_j(t_j) \geq 0$ ; thus

$$R_i^2 = B_i^{1-x_i}(B_*^{1-x_1, 1-x_2}(T)).$$

More generally, action invest is  $(k + 1)$ th-level rationalizable for a type  $t_i$  if he  $(1 - x_i(t_i))$ -believes that  $T$  is  $k$ th-order  $(1 - x_1, 1 - x_2)$ -believed:

$$R_i^{k+1} = B_i^{1-x_i}([B_*^{1-x_1, 1-x_2}]^k(T)).$$

Action invest is rationalizable if it is  $k$ th level rationalizable for all  $k$ . Thus, action invest is rationalizable for both players exactly if  $T$  is common  $(1 - x_1, 1 - x_2)$ -believed:

$$R^\infty = C^{1-x_1, 1-x_2}(T). \quad (2)$$

By a symmetric argument, action Not invest is rationalizable if  $T$  is common  $(x_1, x_2)$ -belief. The next result states this characterization.

**Proposition 1** *Action invest is rationalizable for type  $t_i$  if and only if*

$$t_i \in B_i^{1-x_i}(C^{1-x_1, 1-x_2}(T));$$

*action Not invest is rationalizable for type  $t_i$  if and only if*

$$t_i \in B_i^{x_i}(C^{x_1, x_2}(T)).$$

It is useful to note that, since  $R_i^1 = B_i^{1-x_i}(T)$ ,  $C^{1-x_1, 1-x_2}(T)$  corresponds to a high common belief in the event that action invest is rational. Hence, each part of the proposition states that *an action  $a_i$  is rationalizable for a type  $t_i$  if and only if  $t_i$  assigns sufficiently high probability on a sufficiently high common belief in the event that  $a_i$  is rational*. That is, he finds it sufficiently likely that the action is rational, finds it sufficiently likely that the other player finds it sufficiently likely that the action is rational, . . . , ad infinitum. The key innovation that yields such a simple characterization is taking the threshold for the sufficiency type dependent, depending on the payoff functions of the types throughout.

We report a many-player extension of this characterization in section 4.3.

### 2.3 Risk-Dominant Selection and Multiplicity

Our focus in this paper is on when both actions are rationalizable for both players and when one action is uniquely rationalizable for both players—without loss of generality, we focus on uniqueness of action

invest. Building on the characterization in the previous section, we present intuitive sufficient conditions for each case. Our first result characterizes the cases with multiplicity and uniqueness, as an immediate corollary to the characterization in the previous section.

**Corollary 1** *Both actions are rationalizable for both players if and only if*

$$(t_1, t_2) \in C^{x_1, x_2}(T) \cap C^{1-x_1, 1-x_2}(T).$$

*Invest is the uniquely rationalizable action for a type  $t_i$  if and only if*

$$t_i \in B_i^{1-x_i}(C^{1-x_1, 1-x_2}(T)) \setminus B_i^{x_i}(C^{x_1, x_2}(T)).$$

The first part characterizes the cases with multiple rationalizable solutions. It states that both actions are rationalizable if and only if there is sufficiently high common belief that both actions are rational for both players. The second part characterizes the cases in which invest is the only rationalizable solution. It states that invest is uniquely rationalizable if and only if there is sufficiently high common belief in rationality of invest (i.e.,  $t_i \in B_i^{1-x_i}(C^{1-x_1, 1-x_2}(T))$ ) but there is not sufficiently high common belief in rationality of Not invest (i.e.,  $t_i \notin B_i^{x_i}(C^{x_1, x_2}(T))$ ). Once again, we obtain such simple and straightforward characterizations by taking the threshold for sufficiency type-dependent.

While such characterizations are useful conceptually, they may not be practical. In the rest of this section, we provide simple tractable sufficient conditions for multiplicity and uniqueness. We start with a result for multiplicity:

**Proposition 2** *Both actions are rationalizable whenever there is approximate common certainty that payoffs support multiple strict equilibria, i.e., for some  $\varepsilon \in [0, 1/2]$ ,*

$$(t_1, t_2) \in C^{1-\varepsilon, 1-\varepsilon}(M_\varepsilon)$$

where

$$M_\varepsilon = \{(t'_1, t'_2) \mid \varepsilon \leq x_i(t'_i) \leq 1 - \varepsilon \text{ for both } i\}.$$

*If, in addition,  $x_1$  and  $x_2$  are bounded, then there exist pure-strategy Bayesian Nash equilibria  $s^*$  and  $s^{**}$  on  $T$  such that*

$$s^*(t) = (\text{invest}, \text{invest}) \text{ and } s^{**}(t) = (\text{not invest}, \text{not invest}) \quad (\forall t \in C^{1-\varepsilon, 1-\varepsilon}(M_\varepsilon)).$$

**Proof.** By definition, both actions are 0th-level rationalizable throughout  $C^{1-\varepsilon, 1-\varepsilon}(M_\varepsilon)$ . Towards an induction, assume that both actions are  $k$ th-level rationalizable throughout  $C^{1-\varepsilon, 1-\varepsilon}(M_\varepsilon)$ . Now, for any type  $t_i \in B_i^{1-\varepsilon}(C^{1-\varepsilon, 1-\varepsilon}(M_\varepsilon))$  and any action  $a$ , consider a belief of type  $t_i$  that puts probability 1 on

a  $k$ th-level rationalizable strategy of the other player  $j$  such that each  $t_j \in B_j^{1-\varepsilon}(C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon))$  plays the action  $a$ . Observe that  $\varepsilon \leq x_i(t_i) \leq 1 - \varepsilon$  and that  $t_i$  assigns at least probability  $1 - \varepsilon$  on the other player playing action  $a$ . Hence,  $a$  is a best response to that belief for  $t_i$ , showing that  $a$  is  $(k + 1)$ th-level rationalizable for  $t_i$ . Therefore, both actions are  $(k + 1)$ th-level rationalizable for both players throughout  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$ .

To show the second part, construct an auxiliary game by altering the payoffs of types in  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$  so that they get 1 from invest and 0 from not invest, making invest strictly dominant on  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$ . The auxiliary game satisfies the sufficient conditions of van Zandt (2010) and therefore has a pure strategy Bayesian Nash equilibrium  $s^*$  as in the proposition. But  $s^*$  is also a Bayesian Nash equilibrium in the original game because the types in  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$  play a best response as in the first part of the proof and the other types play a best response by construction. By the same token, we also have a Bayesian Nash equilibrium  $s^{**}$  as in the proposition. ■

Proposition 2 states that both actions are rationalizable whenever there is common  $(1 - \varepsilon, 1 - \varepsilon)$ -belief that both actions would have been  $\varepsilon$ -dominant Nash equilibrium under complete information. This follows a key observation in the robustness literature, going back to Monderer and Samet (1989) which states that any  $p$ -dominant equilibrium of a game can be extended to a larger type space in which the original game is  $(1 - p)$ -evident. In the second part of the proposition, under additional boundedness condition, we use the standard techniques in the robustness literature to construct Bayesian Nash equilibria on  $T$  so that (invest, invest) is played throughout  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$  in one equilibrium and (not invest, not invest) is played throughout  $C^{1-\varepsilon,1-\varepsilon}(M_\varepsilon)$  in another equilibrium. One could use other technical conditions to obtain such results. In the first part, we dispense with those technical conditions by using the weaker solution concept of rationalizability.

We next turn to establishing sufficient conditions under which there is a unique rationalizable action. Our key concept will be approximate uniformity of "rank beliefs", which we now define. We will write  $\bar{r}_i(t_i)$  for the probability that a player assigns to his payoff parameter being greater than or equal to that of the other player, and  $\underline{r}_i(t_i)$  for the probability that it is strictly greater. We will refer to these expressions as "rank beliefs" as they reflect the player's belief about his rank if the players are ordered by the payoff parameter. Formally, we define

$$\begin{aligned}\bar{r}_i(t_i) &= \pi_i(\{t_j | x_j(t_j) \leq x_i(t_i)\} | t_i), \\ \underline{r}_i(t_i) &= \pi_i(\{t_j | x_j(t_j) < x_i(t_i)\} | t_i),\end{aligned}$$

and

$$r_i(t_i) = (\bar{r}_i(t_i) + \underline{r}_i(t_i)) / 2.$$

We refer to  $\bar{r}_i(t_i)$ ,  $\underline{r}_i(t_i)$ , and  $r_i(t_i)$  as the *upper rank belief*, the *lower rank belief* and the *rank belief* of type  $t_i$ , respectively. When  $x_j(\cdot)$  is atomless according to type  $t_i$ , all these beliefs coincide:  $\bar{r}_i(t_i) = \underline{r}_i(t_i) = r_i(t_i)$ . We define upper and lower rank beliefs separately in order to deal with point masses. Such point masses may arise, for example, under complete information.

Rank belief of a type  $t_i$  is uniform if he finds it equally likely that either player's value is higher. Formally, we say that rank belief of a type  $t_i$  is  $\varepsilon$ -uniform if

$$\frac{1}{2} - \varepsilon \leq \underline{r}_i(t_i) \leq \bar{r}_i(t_i) \leq \frac{1}{2} + \varepsilon.$$

We write  $URB_\varepsilon$  for the set of type profiles  $(t_1, t_2)$  where both players have  $\varepsilon$ -uniform rank beliefs:

$$URB_\varepsilon = \left\{ (t_1, t_2) \left| \frac{1}{2} - \varepsilon \leq \underline{r}_i(t_i) \leq \bar{r}_i(t_i) \leq \frac{1}{2} + \varepsilon \text{ for each } i \right. \right\}.$$

We say that rank beliefs are *approximately uniform* if they are  $\varepsilon$ -uniform for some  $\varepsilon \geq 0$ .

Our second concept is a strict version of risk dominance. We say that action invest is  $\varepsilon$ -strictly risk dominant for  $t_i$  if

$$x_i(t_i) \geq \frac{1}{2} + \varepsilon.$$

We write  $SRD_\varepsilon$  for the set of type profiles for which invest is  $\varepsilon$ -strictly risk dominant for each player, so

$$SRD_\varepsilon = \left\{ (t_1, t_2) \left| x_i(t_i) \geq \frac{1}{2} + \varepsilon \text{ for each } i \right. \right\}.$$

We say that action invest is *strictly risk dominant* if invest is  $\varepsilon$ -strictly risk dominant for some  $\varepsilon \geq 0$ .

**Proposition 3** *Invest is the uniquely rationalizable action for both players if it is  $\varepsilon$ -strictly risk dominant for both players and there is common  $(1 - \varepsilon)$ -belief of  $\varepsilon$ -uniform rank beliefs for some  $\varepsilon \geq 0$ , i.e., if*

$$(t_1, t_2) \in SRD_\varepsilon \cap C^{1-\varepsilon, 1-\varepsilon}(URB_\varepsilon).$$

We require only common  $(1 - \varepsilon)$ -belief, not common certainty, of  $\varepsilon$ -uniform rank beliefs for this sufficient condition. But note that in cases where we use this result—to prove results in the fat tails model and to explain the existing global games literature—we actually have common certainty.

Proposition 3 provides a useful sufficient condition for uniqueness, identifying common features of the uniqueness results in the global games literature. It states that invest is uniquely rationalizable if it is strictly risk-dominant and it is common knowledge that the rank beliefs are approximately uniform. Observe that our result establish this result without explicitly assuming some of the critical features of global games, such as existence of dominance regions. Moreover, it allows arbitrary type spaces with minimal continuity and compactness properties. In the rest of this section, we will further discuss our result and present its proof.

Although our game lacks the monotonicity assumptions in global games, we can use rank beliefs to obtain bounds on rationalizable strategies. To this end, we define two cutoffs  $x^*$  and  $x^{**}$  with  $0 \leq x^* \leq x^{**} \leq 1$ :

$$\begin{aligned} x^* &= \inf \{z | z = x_i(t_i) \geq \underline{r}_i(t_i) \text{ for some } i \text{ and } t_i\} \\ x^{**} &= \sup \{z | z = x_i(t_i) \leq \bar{r}_i(t_i) \text{ for some } i \text{ and } t_i\}. \end{aligned}$$

Here,  $x^*$  is the lowest value for which any type of either player has that value and has a lower lower-rank belief; and  $x^{**}$  is the highest value for which any type of any player has that value and a higher upper-rank belief. The next lemma establishes that invest cannot be rationalizable when the value is lower than  $x^*$ , and Not invest cannot be rationalizable when the value exceeds  $x^{**}$ .

**Lemma 3** *Invest is uniquely rationalizable for any  $t_i$  with  $x_i(t_i) > x^{**}$  and Not invest is uniquely rationalizable for any  $t_i$  with  $x_i(t_i) < x^*$ .*

**Proof.** We will show that invest is not rationalizable when  $x_i(t_i) < x^*$ . In particular, let

$$\hat{x}_i \equiv \inf \{x_i(t_i) | t_i \in R_i^\infty\}$$

and assume without loss that

$$\hat{x}_1 \leq \hat{x}_2. \quad (3)$$

By definition, there exists a sequence  $(t_{1,m})$  of types  $t_{1,m} \in R_1^\infty$  such that  $x_1(t_{1,m}) \in [\hat{x}_1, \hat{x}_1 + 1]$  for each  $m$  and  $x_1(t_{1,m}) \rightarrow \hat{x}_1$ . Since  $x_1^{-1}([\hat{x}_1, \hat{x}_1 + 1])$  is sequentially compact, there then exists a convergent subsequence with some limit  $\hat{t}_1$ . Since  $x_1$  is continuous and  $x_1(t_{1,m}) \rightarrow \hat{x}_1$ , we have

$$\hat{x}_1 = x_1(\hat{t}_1). \quad (4)$$

But since  $t_{1,m} \in R_1^\infty$ , we also have  $x_1(t_{1,m}) \geq 1 - \pi_1(R_2^\infty | t_{1,m})$  for all  $m$ . By continuity of  $x_1$  and  $\pi_1$ , we thus have

$$x_1(\hat{t}_1) \geq 1 - \pi_1(R_2^\infty | \hat{t}_1). \quad (5)$$

Therefore,

$$\begin{aligned} \hat{x}_1 &= x_1(\hat{t}_1), \text{ by (4)} \\ &\geq 1 - \pi_1(R_2^\infty | \hat{t}_1), \text{ by (5)} \\ &\geq 1 - \pi_1(\{t_2 | x_2(t_2) \geq \hat{x}_2\} | \hat{t}_1), \text{ by definition of } \hat{x}_2 \\ &\geq 1 - \pi_1(\{t_2 | x_2(t_2) \geq \hat{x}_1\} | \hat{t}_1), \text{ by (3)} \\ &= \pi_1(\{t_1 | x_2(t_2) < x_1(\hat{t}_1)\} | \hat{t}_1) \\ &= \underline{r}_1(\hat{t}_1), \end{aligned}$$

showing that

$$\widehat{x}_2 \geq \widehat{x}_1 \geq x^*.$$

A symmetric argument establishes that not invest is not rationalizable if  $x_i(t_i) > x^{**}$ . ■

When  $x^{**} = 1$  (or  $x^* = 0$ ), Lemma 3 is vacuous, stating that invest is uniquely rationalizable when it is dominant. When  $x^{**} < 1$ , Lemma 3 establishes that invest remains uniquely rationalizable throughout the interval  $(x^{**}, 1)$ .

This is similar to the main result of Carlsson and van Damme (1993). In their result, if one can connect a type  $t_i$  to a type  $\bar{t}_i$  at which invest is a dominant action, via a continuous path along which invest is either risk-dominant or dominant, then invest is uniquely rationalizable at  $t_i$ . Here, we do not explicitly assume the existence of any type with a dominant action and make only minimal assumptions about the structure of the type space. Nonetheless, when  $x^{**} < 1$ , each type  $t_i$  with  $x^{**} < x_i(t_i) < 1$  assigns a substantial probability (i.e. a probability greater than  $1 - x_i(t_i)$ ) on a set of types  $t_j$  whose values are higher than that of  $t_i$  and assign a substantial probability to yet another set of types with similar properties. Under our compactness assumption, this chain leads to types who assign a substantial probability on types for which invest is a dominant action. Such a chain forms a contagion path. Note that the types in the chain have type-dependent thresholds as in Proposition 1.

**Proof of Proposition 3.** Proposition 3 immediately follows from Lemma 3. Under common  $(1 - \varepsilon, 1 - \varepsilon)$ -belief of  $URB_\varepsilon$ , we clearly have  $x^{**} \leq 1/2 + 2\varepsilon$ . Hence, whenever  $(t_1, t_2) \in SRD_\varepsilon \cap C^{1-\varepsilon, 1-\varepsilon}(URB_\varepsilon)$ , we have  $x_i(t_i) > x^{**}$  and invest is uniquely rationalizable (by Lemma 3) for both players. ■

While Proposition 3 will be the focus on our analysis in this paper, to relate our work to the existing literature, we will also note a weaker sufficient condition:

**Corollary 2 (Uniqueness II)** *Invest is the uniquely rationalizable action for both players if it is strictly risk dominant for both and there is approximate common certainty that each player has approximately uniform rank beliefs when his action is strictly risk dominant but not dominant, i.e., for some  $\varepsilon \in [0, 1]$ ,*

$$(t_1, t_2) \subseteq C^{1-\varepsilon, 1-\varepsilon} \left( \left\{ (t'_1, t'_2) \left| \begin{array}{l} x'_i(t'_i) \geq \frac{1}{2} + \varepsilon \text{ and} \\ x'_i(t'_i) \in [\frac{1}{2} + \varepsilon, 1] \Rightarrow \frac{1}{2} - \varepsilon \leq r_i(t'_i) \leq \bar{r}_i(t'_i) \leq \frac{1}{2} + \varepsilon \\ \text{for both } i \end{array} \right. \right\} \right).$$

Note that this sufficient condition merges the uniform rank beliefs and strict risk dominance properties into a single condition. We simply drop the restriction on rank beliefs when a player has a dominant strategy. This will clearly not matter for strategic results.

### 3 One Dimensional Type Space Model

We now restrict our attention to a particular class of symmetric "global game" type spaces. Players' types are one dimensional and drawn according to a common prior. In particular,

$$t_i = \tau\eta + \sigma e_i$$

where  $\tau, \sigma \in (0, 1)$ ;  $\eta$  is a random variable with symmetric, positive, Lipschitz-continuous density function  $g$  (with corresponding cumulative distribution function  $G$ ); and random variables  $(e_1, e_2)$  are independently and identically distributed with symmetric, positive, Lipschitz-continuous density function  $f$  (and corresponding cumulative distribution function  $F$ ). Assume that there exist  $\bar{e} > 0$  and  $\bar{\eta} > 0$  such that  $f$  and  $g$  are monotonically decreasing on  $[\bar{e}, \infty)$  and  $[\bar{\eta}, \infty)$ , respectively. For some  $y \in \mathbb{R}$ , we assume that

$$x_1(t_1) = y + t_1 \text{ and } x_2(t_2) = y + t_2.$$

The functions  $f$  and  $g$ , and the parameters  $y, \sigma$  and  $\tau$  are all common knowledge. Thus each  $(f, g, y, \sigma, \tau)$  describes a type space.

The interpretation of the model is that  $y$  is a "public signal", or prior mean, of the players' types. The noise in their types—i.e., the difference between their values and the prior mean, is the sum of a common shock  $\eta$  and conditionally independent shock  $e_i$ . The size of these common and idiosyncratic shocks are parameterized by  $\tau$  and  $\sigma$  respectively. An equivalent interpretation—in keeping with global games literature—is that there is an unknown "true state"  $\theta$  drawn according to density  $g\left(\frac{\theta-y}{\tau}\right)$  and the players observe signals that are equal to true state plus conditionally independent noise. A player's type is then the difference between this signal of  $\theta$  and prior mean of  $\theta$ .

We will be concerned about players' beliefs about other players' payoffs. We will write  $r_{\sigma,\tau}(t)$  for the probability that type  $t$  assigns to the other player having a lower type, and thus a lower value, than him; so

$$r_{\sigma,\tau}(t) = \int_{\eta=-\infty}^{\infty} F\left(\frac{t-\tau\eta}{\sigma}\right) g(\eta) d\eta.$$

As in the general model, we refer to  $r_{\sigma,\tau}(t)$  as the *rank belief* of  $t$ . Note that  $r_{\sigma,\tau}$  is continuous, as we have a density; also,  $r_{\sigma,\tau}(t)$  is independent of the prior mean  $y$ . (Since we have density, upper and lower rank beliefs coincide.)

In the remainder of this section, we will maintain the assumption that distributions  $f$  and  $g$  have well-behaved "fat tails". In the next section, we will consider the case where  $f$  and  $g$  are standard normal distributions, a special case that has been much studied in the literature.

### 3.1 Fat Tails

In the remainder of this section, we will assume that both common and idiosyncratic components are drawn from fat-tailed distributions. Fat tailed distributions arise naturally in situations of model uncertainty, as discussed in Acemoglu, Chernozhukov, and Wold (2006). When we are interested in approximating complete information, tail properties are central to understanding higher-order beliefs. Assuming away mass at the tails is potentially very misleading. We discuss further issues in using fat tailed distributions following some formal definitions.

A function  $L : \mathbb{R} \rightarrow \mathbb{R}_+$  is said to have *slow-varying tails* if

$$\lim_{x \rightarrow \infty} \frac{L(\lambda x)}{L(x)} = 1$$

for all  $\lambda > 0$ . Convergence is uniform over compact intervals of  $\lambda$ . We assume that  $f$  and  $g$  have *regularly-varying tails*, so that

$$\begin{aligned} f(e) &= AL_f(e) e^{-\alpha} \\ g(\eta) &= BL_g(\eta) \eta^{-\beta} \end{aligned}$$

for some  $\alpha, \beta > 1$ ,  $A, B > 0$ , and for some slowly-varying functions  $L_f$  and  $L_g$ . A particular example of regularly varying distributions is Pareto distribution where  $f$  and  $g$  are constant up to some  $\bar{e}$  and  $\bar{\eta}$ , respectively, and  $f(e)$  and  $g(\eta)$  are proportional to  $e^{-\alpha}$  and  $\eta^{-\beta}$ , respectively, thereafter. We will use Pareto distribution in our numerical examples. Throughout the paper, when we refer to fat-tailed distributions, we are assuming the stronger property that distributions have regularly-varying tails.

By the generalized central limit theorem, the limit of averages can be approximated by a Normal or a Pareto distribution, depending on whether the underlying distribution has finite or infinite variance.

Belief updating under fat tails is quite different from belief updating under normal distribution. Under the normal distribution, a player updates his belief by taking a weighted average of his signals and his prior. This updating is well behaved and gives rise to many monotonicity properties. A player's expectation of his own and the other player's payoff type are increasing in his own signal. A monotone likelihood ratio property is satisfied. And so on.

These properties are not preserved by belief updating under fat tails. However, the structure of updating under fat tails has its own psychological and economic intuition, and we believe it to be very relevant for economic applications. Suppose that a player observes the signal  $t_i = \eta + e_i$ . What is his expectation of the signal of the other player  $t_j = \eta + e_j$ . If the signal  $t_i$  is close to its prior mean of 0, the player's expectation of  $\eta$ , and thus of  $t_j$ , will be—roughly speaking—an average of his prior 0 and his signal  $t_i$ , as in the normal case. But if signal  $t_i$  is a long way from 0, the player will think it most likely that either  $\eta$  or  $e_i$ , but not both, were different from 0. Which one he thinks most likely depends on the shape of the tails.






Figure 1: Expectation of the other player's type conditional on ones own when the idiosyncratic components have thicker tails than the common component (i.e.,  $\alpha < \beta$ ).

This is illustrated in Figure 1 for the case of Pareto distributions, where the  $e_i$  are drawn from the Pareto distribution with tail parameter  $\beta = 4$  and  $\eta$  is drawn from a Pareto distribution with a thinner tail  $\alpha = 2$ . For small positive  $t_i$ , he updates his expectation of  $\eta$  and thus his expectation of the other player's type  $t_j$ . So his expectation of  $t_j$  is increasing in  $t_i$  for a while. But if he observes a substantially larger  $t_i$ , he then questions whether his private signal is valuable, attributing a substantial part of this deviation to  $e_i$ . This leads to a lower expectation about  $t_j$  than the expectation he would have after observing a lower signal  $t_i$  (as in the decreasing part of the graph). In the limit, when he observes an extremely positive signal, he discredits his private signal altogether as a meaningless noise and sticks to his prior, expecting  $E[t_j | t_i] = 0$ . This is a behaviorally intuitive form of updating which is also rational. While such non-monotone behavior prevents us from using many existing techniques in the global games literature, the above updating behavior under regularly-varying tails provide us other means that lead to sharper characterizations and economically natural results.

### 3.2 Higher-Order Beliefs

A key property of updating with fat tailed distributions is that when a type  $t_i$  is in the tail, he will attribute all the deviation from the mean to the shock with thicker tails. When the idiosyncratic shock has thicker tails ( $\beta > \alpha$ ), as in the above example, he will attribute the deviation to the idiosyncratic shock and will believe that the other player's type is near the origin. This will lead to approximate common certainty of the payoffs and thereby to multiplicity. On the other hand, when the common shock has thicker tails ( $\alpha > \beta$ ), he will attribute the deviation to the common shock and will have approximately uniform rank beliefs, leading to uniqueness. We start with our key result for this section, which applies in the latter case

and applies for all levels of noise. We then consider limiting results as noise goes to zero and thus as we approach complete information.

### 3.2.1 Thinner Idiosyncratic Tails

Our main result says that if the idiosyncratic shock (or private information) has thinner tails than the common shock (or public information) and the idiosyncratic term is of sufficiently smaller size, then there is common certainty of approximately uniform rank beliefs.

**Proposition 4** *If  $\alpha > \beta + 1$  and  $\sigma/\tau$  is sufficiently small, then there is common certainty of approximately uniform rank beliefs, i.e., for any  $\varepsilon > 0$ , there exists  $\bar{\xi} > 0$  such that whenever  $\sigma/\tau < \bar{\xi}$ ,*

$$\frac{1}{2} - \varepsilon < r_{\sigma,\tau}(t) < \frac{1}{2} + \varepsilon \quad (\forall t).$$

Note that the requirement on the size of the shocks is that  $\sigma/\tau$  is smaller than a bound  $\bar{\xi}$ , which depends only on  $\varepsilon$ ,  $f$ , and  $g$ . This result does not require  $\sigma$  and  $\tau$  to be small and thus gives a sufficient condition for common certainty of approximately uniform rank beliefs away from the complete information limit.

For the types in the tail, because the idiosyncratic term has a thinner tail, a player attributes the aberration to the common shock and believes that other player's type is near his own, having an approximately uniform rank belief. For the types away from the tail, since the idiosyncratic term is of sufficiently smaller size, he again attributes the deviation from the mean to the common shock and has an approximately uniform rank belief. This is vividly illustrated in Figure 2, where  $\alpha = 4$  and  $\beta = 2$ . Since  $\alpha > \beta$ , for all values of  $(\sigma, \tau)$ , the rank beliefs  $r_{\sigma,\tau}(t)$  approach  $1/2$  near the tail. Away from the tail,  $r_{\sigma,\tau}(t)$  may be far from  $1/2$  when  $\sigma/\tau$  is large, as in the case of  $(\sigma, \tau) = (1, 1)$ . But when  $\sigma/\tau$  is small,  $r_{\sigma,\tau}$  is approximately  $1/2$  everywhere. Therefore, we have common certainty of approximately uniform rank beliefs. Our proof formalizes these ideas next.

**Proof.** Writing  $\xi = \sigma/\tau$ , note that  $r_{\sigma,\tau}(t) = r_{\xi,1}(t/\tau)$ . Hence, it suffices to show that there exists  $\bar{\xi}$  such that  $r_{\xi,1}(t) < 1/2 + \varepsilon$  for all  $t \geq 0$  whenever  $\xi < \bar{\xi}$ ; throughout the proof, we will assume that  $\xi \ll 1/2$  is sufficiently small. All the other bounds follow from symmetry. To this end, we first observe that, since  $g$  has regularly varying tails, there exists  $t_0$  such that for all  $t > t' \geq t_0$ ,

$$(t/t')^{-\beta-1} / (1 + \varepsilon/2) \leq \frac{g(t)}{g(t')} \leq (1 + \varepsilon/2) (t/t')^{-\beta+1}. \quad (6)$$

We write

$$\tilde{e}(t) = \max \{t, \delta/\xi^\gamma\} \geq \bar{e} \quad (7)$$

for some large  $\delta$  and for some  $\gamma \in (1/(\alpha - \beta), 1)$ . We suppress the dependence on  $t$  in the sequel. We observe that

$$r_{y,\xi,1}(t) \leq (I_1 + I_2) / I_3$$




Figure 2: Rank beliefs for  $\sigma = 1, 0.1, 0.001$  and  $(\alpha, \beta, \tau) = (4, 2, 1)$ .

where

$$\begin{aligned}
 I_1 &= \int_{-\xi\tilde{e}}^{\xi\tilde{e}} f(z/\xi) F(z/\xi) g(t-z) dz \leq \frac{1}{2} (F(\tilde{e}) - F(-\tilde{e})) \xi g^{M,\xi}(t), \\
 I_2 &= \int_{z \notin (-\xi\tilde{e}, \xi\tilde{e})} f(z/\xi) F(z/\xi) g(t-z) dz \leq f(\tilde{e}), \\
 I_3 &= \int_{-\xi\tilde{e}}^{\xi\tilde{e}} f(z/\xi) g(t-z) dz \geq (F(\tilde{e}) - F(-\tilde{e})) \xi g^{m,\xi}(t)
 \end{aligned}$$

and

$$g^{M,\xi}(t) = \max_{z \in [t-\xi\tilde{e}, t+\xi\tilde{e}]} g(t) \text{ and } g^{m,\xi}(t) = \min_{z \in [t-\xi\tilde{e}, t+\xi\tilde{e}]} g(t).$$

Combining the above inequalities, we conclude that

$$r_{y,\xi,1}(t) \leq \frac{1}{2} \frac{g^{M,\xi}(t)}{g^{m,\xi}(t)} + \frac{f(\tilde{e})}{(F(\tilde{e}) - F(-\tilde{e})) \xi g^{m,\xi}(t)}. \tag{8}$$

Now, note that, since  $\alpha > \beta + 1$ , there exists  $\delta$  such that

$$\frac{f(z/\xi^\gamma)}{(F(z/\xi^\gamma) - F(-z/\xi^\gamma)) \xi g(2z/\xi^\gamma)} < \varepsilon/2 \quad \forall z \geq \delta.$$

We pick such a  $\delta$  in definition of  $\tilde{e}$ . Then, using the fact that  $g^{m,\xi}(t) \geq g(t + \tilde{e}(t)) \geq g(2\tilde{e}(t))$ , one can easily check that

$$\frac{f(\tilde{e}(t))}{(F(\tilde{e}(t)) - F(-\tilde{e}(t))) \xi g^{m,\xi}(t)} < \varepsilon/2 \quad \forall t. \tag{9}$$

In order to find an upper bound for  $\frac{1}{2}g^{M,\xi}(t)/g^{m,\xi}(t)$ , consider only  $\xi \ll (t_0/\delta)^{1/(1-\gamma)}$ , so that  $\tilde{e}(t_0) = t_0$ . Then, for any  $t > t_0/(1-\xi)$ , we have  $t - \xi\tilde{e} = (1-\xi)t > t_0$ , and hence, by (6),

$$\frac{g^{M,\xi}(t)}{g^{m,\xi}(t)} \leq (1 + \varepsilon/2) \left( \frac{t + \xi\tilde{e}}{t - \xi\tilde{e}} \right)^{\beta+1} = (1 + \varepsilon/2) \left( \frac{1 + \xi}{1 - \xi} \right)^{\beta+1}. \quad (10)$$

On the other hand, since  $g$  is Lipschitz continuous and positive, there also exists  $\kappa > 0$  such that

$$g(t') \leq g(t) (1 + (t' - t)\kappa)$$

for any  $t, t' \in [0, 2t_0]$ . (Note that  $\kappa$  is independent of  $\xi$ .) Hence, for any  $t \in [0, t_0/(1-\xi)]$ , we also have

$$\frac{g^{M,\xi}(t)}{g^{m,\xi}(t)} \leq \frac{1 + \kappa\xi\tilde{e}}{1 - \kappa\xi\tilde{e}} \leq \frac{1 + \kappa \max\{\xi t_0/(1-\xi), \xi^{1-\gamma}\delta\}}{1 - \kappa \max\{\xi t_0/(1-\xi), \xi^{1-\gamma}\delta\}}. \quad (11)$$

Substituting (9-11) in (8), we obtain

$$r_{y,\xi,1}(t) \leq \frac{1}{2} \max \left\{ \frac{1 + \kappa \max\{\xi t_0/(1-\xi), \xi^{1-\gamma}\delta\}}{1 - \kappa \max\{\xi t_0/(1-\xi), \xi^{1-\gamma}\delta\}}, (1 + \varepsilon/2) \left( \frac{1 + \xi}{1 - \xi} \right)^{\beta+1} \right\} + \varepsilon/2.$$

Observe that the right-hand side is independent of  $t$  and converges to  $1/2 + 3\varepsilon/4$  as  $\xi \rightarrow 0$ . Therefore, there exists  $\bar{\xi} > 0$  such that for all  $\xi \in (0, \bar{\xi})$ ,

$$r_{y,\xi,1}(t) < \frac{1}{2}(1 + \varepsilon) + \varepsilon/2 = \frac{1}{2} + \varepsilon \quad (\forall t \geq 0, \forall \xi < \bar{\xi}).$$

■

### 3.2.2 Approximating Complete Information

We now report results for when  $\sigma$  and  $\tau$  are both small, analyzing what happens to higher-order beliefs along sequences  $(\sigma_n, \tau_n) \rightarrow (0, 0)$ . Thus we will be considering a sequences where there is very little uncertainty about payoffs and, in particular, types converge in the product topology to a complete information model, where there is common certainty of payoffs.

For any given  $(\sigma_n, \tau_n)$ , we can consider the beliefs of a player observing signal  $t = \tau_n\eta + \sigma_n e$  about the common component of his signal  $\tau_n\eta$ . Belief about other variables will then be derived from this. We write  $\mu_n(\cdot|t)$  for the probability distribution of  $\tau_n\eta$  conditional on  $(\sigma_n, \tau_n, t)$ .<sup>5</sup> We are interested in characterizing the limit of  $\mu_n(\cdot|t)$  as  $n \rightarrow \infty$ .

<sup>5</sup>The probability of any interval  $(\tau_n\eta_1, \tau_n\eta_2)$  under  $\mu_n(\cdot|t)$  is

$$\mu_n((\tau_n\eta_1, \tau_n\eta_2)|t) = I_n(t - \tau_n\eta_1, t - \tau_n\eta_2) / I_n(-\infty, \infty)$$

where

$$I_n(z_1, z_2) = \int_{z_1}^{z_2} f(z/\sigma_n) g\left(\frac{t-z}{\tau_n}\right) dz.$$

Towards this goal, we define

$$q_n(t) = \frac{\sigma_n g(t/\tau_n)}{\sigma_n g(t/\tau_n) + \tau_n f(t/\sigma_n)},$$

and assume that  $q_n(t)$  converges to some  $q(t) \in [0, 1]$  for each  $t$ , and that the convergence is uniform over  $[\underline{t}, \bar{t}]$  for any  $\bar{t} > \underline{t} > 0$ . Under regularly-varying tails, it is straightforward to show that

$$q_n(t) \cong \frac{1}{1 + \rho_n t^{\beta-\alpha} A/B} \text{ and } q(t) = \frac{1}{1 + \rho t^{\beta-\alpha} A/B}$$

where

$$\rho_n = \frac{\sigma_n^{\alpha-1} L_f(1/\sigma_n)}{\tau_n^{\beta-1} L_g(1/\tau_n)} \text{ and } \rho = \lim_{n \rightarrow \infty} \rho_n.$$

In the case of Pareto distribution, we simply have  $\rho_n(t) = \sigma_n^{\alpha-1}/\tau_n^{\beta-1}$  allowing a tractable analysis based on the tail indices  $\alpha$  and  $\beta$  of idiosyncratic and common components, respectively. In general, except for knife-edge cases,  $\rho$  is either 0 or  $\infty$ , and  $q(t)$  is either 1 or 0, independent of  $t$ . Finally, we define our notion of convergence for type-dependent probability distributions as follows. For any sequence of type-dependent probability distributions  $P_n(\cdot|\cdot)$  and any  $P(\cdot|\cdot)$ , where  $P_n(\cdot|t_i)$  and  $P(\cdot|t_i)$  are probability distributions over the same type space for each  $t_i$ , we say that  $P_n(\cdot|\cdot)$  converges to  $P(\cdot|\cdot)$  uniformly over compact sets of types if  $\int h dP_n(\cdot|t_i) \rightarrow \int h dP(\cdot|t_i)$  uniformly over types  $t_i \in [\underline{t}, \bar{t}]$  for any bounded continuous function  $h$  and any compact interval  $[\underline{t}, \bar{t}]$ .

The following result establishes the implication of fat tails that, in the limit, players rely only on one of the public and the private signals, ignoring the other.

**Lemma 4** *As  $n \rightarrow \infty$ , the beliefs  $\mu_n(\cdot|\cdot)$  about the common shock  $\tau_n \eta$  converge to  $\mu_\infty(\cdot|\cdot)$  uniformly over compact sets of types where*

$$\mu_\infty(0|t_i) = 1 - q(t_i) \text{ and } \mu_\infty(t_i|t_i) = q(t_i)$$

for each  $t_i \neq 0$  and  $\mu_\infty(0|0) = 1$ .

Thus, in the limit  $(\sigma_n, \tau_n) \rightarrow (0, 0)$ , each individual puts positive probability only on two points, entertaining only two possible theories. He puts probability  $q(t_i)$  on  $(\eta = 0, \sigma_n e_i = t_i)$ , attributing all of the variation in his type  $t_i$  from 0 to the noise  $e_i$  in his individual information. He puts probability  $1 - q(t_i)$  on  $(\tau_n \eta = t_i, e_i = 0)$ , attributing all of the variation in his type  $t_i$  from 0 to the common shock. Consequently, the players have stark limiting beliefs regarding their relative ranking. When the tails of the idiosyncratic components are fatter than that of the common component ( $\alpha < \beta$ ), we have  $q(t_i) = 1$ , and he ends up attributing all the deviation to his individual information, maintaining the belief that the other player's type is around 0, as in (the extremes of) Figure 1. When the tails of the idiosyncratic components

are thinner than that of the common component ( $\alpha > \beta$ ), we have  $q(t_i) = 0$ , and he ends up attributing all the deviation to the common shock, maintaining the belief that the other player's type is around his own, as in (the extremes of) Figure 3 below.

Identifying the limiting rank beliefs requires more care because, conditional on  $t_i$ , the limit beliefs regarding  $t_j$  are somewhat complicated. In the limit, by Lemma 4, player  $i$  assigns probability  $1 - q(t_i)$  on  $\tau\eta = 0$  and probability  $q(t_i)$  on  $\tau\eta = t_i$ . But in the limit case of  $\tau\eta = t_j = t_i$ , the relative ranking of  $t_j$  and  $t_i$  are not determined from Lemma 4 because the limit CDF of  $t_j$  conditional on  $\tau\eta$  is discontinuous at  $\tau\eta$ .<sup>6</sup> After all, probability of  $t_j = t_i$  is zero. The next result establishes the limiting rank beliefs, defined as

$$r_\infty(t_i) \equiv \lim_{n \rightarrow \infty} r_{\sigma_n, \tau_n}(t_i) = \lim_{n \rightarrow \infty} \Pr(t_j \leq t_i | t_i, \sigma_n, \tau_n).$$

The result shows that when the noise is small so that  $t_i$  is near  $\tau\eta$ , the player assigns probability 1/2 on the other player's signal being lower than his own.

**Lemma 5** *For any  $\bar{t} > \underline{t} > 0$ , as  $n \rightarrow \infty$ ,  $r_{\sigma_n, \tau_n}(t_i)$  uniformly converges to*

$$r_\infty(t) = \begin{cases} 1 - q(t)/2 & \text{if } t > 0 \\ q(t)/2 & \text{if } t < 0 \end{cases}$$

over  $[-\bar{t}, -\underline{t}] \cup [\underline{t}, \bar{t}]$ .

The limiting rank-beliefs have an interesting form. When  $\rho \equiv \lim_{n \rightarrow \infty} \frac{\sigma_n^{\alpha-1} L_f(1/\sigma_n)}{\tau_n^{\beta-1} L_g(1/\tau_n)} = \infty$ , we have  $q(t) \equiv 0$ , and the limiting rank belief is a step function at  $t = 0$ . The player nearly attributes all of the variation in his signal to the noise in his own private information and keeps believing that  $t_j$  is around zero—and that  $\theta$  and  $x_j$  are around  $y$ . Since both players do so for all types, it is then easy to obtain a high common belief in the event that both players assign high probability on  $\theta$  being around  $y$ . The next corollary establishes this formally.

**Corollary 3** *If  $\rho = \infty$  and  $\underline{x} < y < \bar{x}$ , then, whenever payoffs are in  $[\underline{x}, \bar{x}]$ , there is limit approximate common certainty that payoffs are in  $[\underline{x}, \bar{x}]$ , i.e., for any  $\varepsilon > 0$ , there exists  $\bar{n}$  such that, for all  $n \geq \bar{n}$ ,*

$$[\underline{x} - y, \bar{x} - y]^2 \subseteq C^{1-\varepsilon, 1-\varepsilon} \left( [\underline{x} - y, \bar{x} - y]^2 \right)$$

in the  $(\sigma_n, \tau_n)$  type space, where  $[\underline{x} - y, \bar{x} - y]^2$  is the set of type profiles under which the payoffs are in  $[\underline{x}, \bar{x}]$ .

**Proof.** Since  $\mu_n(\cdot | t_i) \rightarrow \mu_\infty(\cdot | t_i)$  uniformly over  $t_i \in [\underline{x} - y, \bar{x} - y]$  (by Lemma 4), and since  $\sigma_n \rightarrow 0$ , there exists  $\bar{n}$  such that, for all  $n \geq \bar{n}$  and for all  $t_i \in [\underline{x} - y, \bar{x} - y]$ , the probability that  $t_j \equiv \tau_n \eta + \sigma_n e_j \in$

<sup>6</sup>With our formulation, the rank beliefs is 1/2 in the limit.




Figure 3: Rank beliefs under  $\alpha = 4 > 2 = \beta$  for  $\sigma = \tau = 1$ ,  $\sigma = \tau = 0.1$ , and  $\sigma = \tau = 0.01$ .

$[\underline{x} - y, \bar{x} - y]$  is at least  $\mu_\infty(0|t_i) - \varepsilon$ . But since  $\rho = \infty$ , we have  $\mu_\infty(0|t_i) = 1$ . Therefore, for all  $n \geq \bar{n}$ , all types  $t_i \in [\underline{x} - y, \bar{x} - y]$  assign at least probability  $1 - \varepsilon$  on  $t_j \in [\underline{x} - y, \bar{x} - y]$ , as desired. ■

That is, when the common component has thinner tails than the idiosyncratic component (e.g.,  $\alpha < \beta$  and  $\sigma_n/\tau_n \gg 0$  so that  $\rho = \infty$ ), the event that the payoffs are around their ex-ante mean is a  $(1 - \varepsilon, 1 - \varepsilon)$ -evident event: whenever it happens, the players assign at least probability  $1 - \varepsilon$  on that event. This will further imply that the equilibria of the complete information game extend to the  $(\sigma_n, \tau_n)$  type space.

On the other hand, if  $\alpha > \beta$  and  $\sigma_n/\tau_n$  is bounded so that the common component has fatter tails than the idiosyncratic component, then  $\rho = 0$  and  $q(t_i) = 1$  everywhere. Our lemma then implies that the rank beliefs converge to  $r_\infty(t_i) = \frac{1}{2}$  on compact intervals of  $t_i$  that exclude 0. However, this does not deliver common certainty of approximately uniform rank beliefs, because we do not have uniform convergence around zero. The problem is vividly illustrated in Figure 3. Although  $r_{\sigma,\tau}$  converges to  $1/2$  pointwise everywhere,  $r_{\sigma,\tau}$  traces the entire interval  $[0.2, 0.8]$  around  $t_i = 0$  no matter how small  $(\sigma, \tau)$  is. (This is because when one fixes the ratio of  $\sigma/\tau$ , reducing the noise is equivalent to shrinking  $t_i$  around 0, i.e.,  $r_{a\sigma, a\tau}(t_i) = r_{\sigma,\tau}(t_i/a)$ .) To obtain common certainty of approximately uniform rank beliefs, we need approximately uniform rank beliefs for types around 0, and for this it is necessary to have  $\sigma$  small relative to  $\tau$ . However, Proposition 4 already established this above—for both large and small  $(\sigma, \tau)$ . In that case,  $r_{\sigma,\tau}$  converges to  $1/2$  uniformly everywhere as  $\sigma/\tau$  gets small (see Figure 2). This leads to common-certainty of approximate rank beliefs in the limit:

**Corollary 4** *Assume  $\alpha > \beta + 1$  and  $\lim_{n \rightarrow \infty} \frac{\sigma_n}{\tau_n} = 0$ . Then, there is limit common certainty of approxi-*

mately uniform rank beliefs, i.e., for any  $\varepsilon > 0$ , there exists  $\bar{n}$  such that if  $n \geq \bar{n}$ ,

$$\frac{1}{2} - \varepsilon < r_{\sigma_n, \tau_n}(t) < \frac{1}{2} + \varepsilon \quad (\forall t)$$

in the  $(\sigma_n, \tau_n)$  type space.

### 3.3 Multiplicity and Uniqueness of Rationalizable Actions

We can now immediately apply our results about higher-order beliefs in the fat-tails model to the general strategic analysis in Section 2.2 to derive strategic results in the fat-tails model.

The sufficient conditions for common certainty of uniform rank beliefs in Proposition 4 immediately imply the selection of the risk-dominant action as the unique rationalizable action, as established by our general result Proposition 3:

**Corollary 5** *If  $\alpha > \beta + 1$ , invest is strictly risk dominant for both players and  $\sigma/\tau$  is sufficiently small, then invest is uniquely rationalizable for both players, i.e., for any  $\varepsilon > 0$ , there exists  $\bar{\xi} > 0$  such that whenever  $\sigma/\tau < \bar{\xi}$ , if  $x_i(t_i) > \frac{1}{2} + \varepsilon$  for both  $i$ , invest is uniquely rationalizable for both players.*

We can also use the higher-order belief results about approximately complete information to establish conditions for "limit multiplicity"—i.e., multiple rationalizable actions as we approach complete information—or "limit uniqueness". In particular, we conclude that there is multiplicity whenever the common component has a thinner tail than the idiosyncratic components, and conversely, the risk-dominant action is uniquely rationalizable whenever the idiosyncratic components have thinner tails. Building on Proposition 2 and Corollary 3, our first result establishes limit multiplicity in the former case.

**Corollary 6** *If  $(\sigma_n, \tau_n) \rightarrow (0, 0)$  and  $\rho = \lim_{n \rightarrow \infty} \frac{\sigma_n^{\alpha-1} L_f(1/\sigma_n)}{\tau_n^{\beta-1} L_g(1/\tau_n)} = \infty$ , then there is limit multiplicity, i.e., for any  $0 < y < 1$  and  $\varepsilon > 0$ , there exists  $\bar{n}$  such that, in every  $(\sigma_n, \tau_n)$  type space with  $n \geq \bar{n}$ , both actions are rationalizable for both players whenever  $x_i(t_i) \in [\varepsilon, 1 - \varepsilon]$  for both  $i$ .*

**Proof.** Let  $\rho = \infty$  and take any  $\varepsilon > 0$ . Assume without loss of generality that  $\varepsilon < y < 1 - \varepsilon$ . Then, Corollary 3 establishes that the event  $M_\varepsilon$  that the payoffs are within the interval  $[\varepsilon, 1 - \varepsilon]$  is  $(1 - \varepsilon, 1 - \varepsilon)$ -evident whenever  $n$  is sufficiently large. Then, for any such  $n$ , Proposition 2 establishes that both actions are rationalizable throughout  $M_\varepsilon$ . ■

The multiplicity here is illustrated in Figure 4. When  $(\sigma, \tau) = (0.1, 0.1)$ , the rank beliefs are as in the flatter curve. For  $y = 1/2$ , this curve intersect the value function  $x_i$  at three points when  $y = 0.5$ , one at  $x^* \simeq 0$ , one at  $x^{**} \simeq 1$ , and one in the middle. Each of the extremal intersections corresponds to a symmetric equilibrium. In one equilibrium, players invest iff their values exceed  $x^*$ , and in another equilibrium they invest iff their values exceed  $x^{**}$ , yielding multiple solutions for types with values in






Figure 4: The rank beliefs for  $(\sigma, \tau) = (0.1, 0.1)$  and  $(\sigma, \tau) = (0.01, 0.01)$  under  $(\alpha, \beta) = (2, 4)$ . The straight lines corresponds to  $x_i$  for  $y = 0, 0.2, 0.5, 0.8, 1$ .

$(x^*, x^{**})$  as in the corollary. For this value of  $(\sigma, \tau)$ , such multiplicity is supported for an interval  $(\underline{y}, \bar{y})$ , where  $\underline{y}$  is slightly above 0.2 and  $\bar{y}$  is slightly below 0.8. For values  $y = 0.2$  and  $y = 0.8$ , we have a unique rationalizable action everywhere except for the type corresponding to the unique intersection (by Lemma 3). When the noise reduces to  $(\sigma, \tau) = (0.1, 0.1)$ , the rank beliefs become as in the steeper curve. Now, a larger set values for  $y$  leads to multiple equilibria, including  $y = 0.2$  and  $y = 0.8$ , as the corresponding value functions intersect the new curve at multiple points. In the limit  $(\sigma, \tau) \rightarrow (0, 0)$ , the curve becomes the step function at  $t_i = 0$ , yielding multiple equilibria for all values of  $y \in (0, 1)$ .

The key insight for this result is as follows. When the players' private information is weaker than the public information, each player attributes all of the deviation in his type from the ex-ante mean 0 to the noise in his private information and keeps believing that the other player's payoff is near the ex-ante mean  $y$ . Since this is true for all types of both players, such high confidence is maintained throughout the players' higher-order beliefs, resulting in a high common belief that the payoffs are near the ex-ante mean  $y$  whenever this is indeed the case. Then, the rationalizable behavior under the complete information about payoffs around  $y$  extend to the case of incomplete information with "approximate common knowledge" about the payoffs. Since both actions are rationalizable under complete information, both actions remain rationalizable under the latter case of incomplete information.

When the players' private information is stronger than their public information (e.g. the tail index  $\alpha$  of the idiosyncratic component is larger than the tail index  $\beta$  of the common component and  $\sigma_n/\tau_n$  is bounded), the opposite happens. Now, each player attributes all of the deviation in his type from the ex-ante mean 0 to the noise in the public information and comes to believe that the other players' payoff is similar to his own, having equal probability of being higher or lower than his own. In such a model, one

cannot maintain any high level of common belief regarding the *whereabouts of the payoffs*. Instead, one maintains a common certainty about the players' rank beliefs being approximately  $1/2$ , as established by Proposition 4. But, such common certainty of the rank beliefs leads to selection of the risk-dominant action as the unique rationalizable action, as established by our general result Proposition 3. Since Proposition 4 applies more generally, such a uniqueness is maintained away from the limit as well as in the limit:

**Corollary 7** *If  $(\sigma_n, \tau_n) \rightarrow (0, 0)$ ,  $\lim_{n \rightarrow \infty} \frac{\sigma_n}{\tau_n} = 0$  and  $\alpha > \beta + 1$ , then there is limit uniqueness, i.e., for any  $\varepsilon > 0$ , there exists  $\bar{n}$  such that, if  $n \geq \bar{n}$ , invest is uniquely rationalizable for both players if  $x_i(t_i) \geq \frac{1}{2} + \varepsilon$  for both  $i$ , in the  $(\sigma_n, \tau_n)$  type space.*

The unique selection above is illustrated in Figure 5. As  $\sigma/\tau$  decreases, the rank belief  $r_{\sigma, \tau}$  uniformly converges to  $1/2$ , restricting the  $x$  values corresponding to an intersection of the rank belief  $r_{\sigma, \tau}$  and  $x_i$  to a smaller  $\varepsilon$  neighborhood of  $1/2$ . This ensures the unique selection of strictly risk-dominant actions, *uniformly* over the possible payoff functions  $x_i$  and the sizes of the idiosyncratic and common shock ( $\sigma$  and  $\tau$ ). For example, the selection here does not depend on the value of  $y$ .

Several remarks are in order. First, there can be multiple equilibria for a given payoff function; for example,  $r_{1,1}$  intersects  $x_i = t_i + 1/2$  at three points, but the intersections are all contained in  $(-0.2, 0.2)$ , allowing the selection of 0.2-strict risk dominant actions. Second, one can switch back and forth between multiplicity and uniqueness as  $\sigma/\tau$  decreases; for example, we have multiplicity under  $(\sigma, \tau) = (1, 1)$  but unique selection under  $(\sigma, \tau) \in \{(2, 1), (0.1, 1), (0.01, 1)\}$ . Finally, under any  $(\sigma, \tau)$  with unique selection, we could still have multiple equilibria by considering another payoff function, e.g.,  $x_i = at_i + 1/2$  for sufficiently small  $a$ , allowing a wide range of types with multiple actions. But all those types will also have values that are near the risk-dominance cutoff  $1/2$ , and we will ultimately be able to select strictly risk dominant action as the unique rationalizable solution whenever such an action exists.

## 4 Discussion: Extensions and Relation to The Literature

### 4.1 Thin Tails and The Normal Case

We restricted our analysis in section 3 to the case of fat tailed distributions. What happens if we allowed thin tailed distributions - where tail densities converge to zero at an exponential rate - in the one dimensional signal setting of section 3.

Some extensions are straightforward. If the common component has a fat tailed distribution and the idiosyncratic component has a thin tailed distribution, then there will be common certainty of uniform rank beliefs in the limit. Conversely, if the common component has a thin tailed distribution and the idiosyncratic component has a fat tailed distribution, then there will be approximate common certainty




Figure 5: Rank beliefs for  $\sigma = 2, 1, 0.1,$  and  $0.01$  under  $(\alpha, \beta, \tau) = (4, 2, 1)$ ; the maximum deviation from  $1/2$  is decreasing in  $\sigma$ . The straight lines correspond to  $x_i$  for  $y = 0.2, .5, 0.8$ .

of payoffs in the limit. Of course, a distribution with bounded support is a special case of a thin tailed distribution.

We do not have general results for the case where both distributions have thin tails. However, a leading case from the global games literature corresponds to assuming a particular thin-tailed distribution - the normal distribution - for both common and idiosyncratic shocks. In this sub-section, we will review the well known analysis of this case to highlight how it fits into the framework of this paper.

Thus we will consider the one dimensional model in the previous section but under the assumption that  $\eta$  and  $e_i$  have standard normal distributions and demonstrate how known results for this problem are implied by the common belief foundations described above. In this case, rank beliefs can be characterized in closed form, and this characterization has been extensively used in the existing literature, e.g., Morris and Shin (2001) and Morris and Shin (2003a).

Player  $i$  with type  $t_i = \tau\eta + \sigma e_i$  will believe that  $\tau\eta$  is distributed with mean

$$\frac{\tau^2}{\sigma^2 + \tau^2} t_i$$

and variance

$$\frac{\sigma^2 \tau^2}{\sigma^2 + \tau^2}$$

and so he will believe that  $t_j = \tau\eta + \sigma e_j$  is distributed with mean

$$\frac{\tau^2}{\sigma^2 + \tau^2} t_i$$

and variance

$$\frac{\sigma^2 \tau^2}{\sigma^2 + \tau^2} + \sigma^2 = \frac{\sigma^2 (\sigma^2 + 2\tau^2)}{\sigma^2 + \tau^2}.$$

Thus the player assigns probability

$$\Phi \left( \sqrt{\frac{\sigma^2 + \tau^2}{\sigma^2 (\sigma^2 + 2\tau^2)}} \frac{\sigma^2}{\sigma^2 + \tau^2} t_i \right) = \Phi \left( \sqrt{\frac{\sigma^2}{(\sigma^2 + 2\tau^2) (\sigma^2 + \tau^2)}} t_i \right)$$

to the other player having a lower signal. Thus the probability that a player with type  $t$  has the higher type is

$$r_{\sigma, \tau}(t) = \Phi \left( \sqrt{\frac{\sigma^2}{(\sigma^2 + 2\tau^2) (\sigma^2 + \tau^2)}} t \right).$$

What can we say about higher-order beliefs in this case? Observe that, as  $\sigma_n \rightarrow 0$  and  $\tau_n \rightarrow 0$ ,  $\sqrt{\frac{\sigma_n^2}{(\sigma_n^2 + 2\tau_n^2)(\sigma_n^2 + \tau_n^2)}}$  converges to  $\infty$  if  $\frac{\sigma_n}{\tau_n} \rightarrow \infty$  and converges to 0 if  $\frac{\sigma_n}{\tau_n} \rightarrow 0$ . Thus if  $\frac{\sigma_n}{\tau_n} \rightarrow \infty$ , we obtain approximate common certainty of payoffs and we have limit multiplicity as defined in Definition ???. If  $\frac{\sigma_n}{\tau_n} \rightarrow 0$ , we do have pointwise convergence to uniform rank beliefs, so that  $r_{\sigma_n, \tau_n}(t) \rightarrow \frac{1}{2}$  for each  $t$ . But this convergence is not uniform in the tails, so that for any  $n$ ,  $r_{\sigma_n, \tau_n}(t) \rightarrow 1$  as  $t \rightarrow \infty$  and  $r_{\sigma_n, \tau_n}(t) \rightarrow 0$  as  $t \rightarrow -\infty$ . Thus there is never approximate common certainty of approximately uniform rank beliefs. But for any  $\varepsilon$ , if we look at the set of  $t$  where rank beliefs are within  $\varepsilon$  of  $\frac{1}{2}$ , we have—by construction—that the marginal type puts a high probability (greater than  $\frac{1}{2} - \varepsilon$ ) to his opponent having a type outside the set. However, it is enough for global game results that there is common certainty that rank beliefs are close  $\frac{1}{2}$  when players do not have a dominant strategy. Thus, the weaker sufficient condition of Lemma 2 for limit uniqueness is satisfied and we have limit uniqueness as defined in Definition ???.

Thus we find that the fat tailed limit results deliver sharper sufficient conditions for limit uniqueness while the well known normal results achieve limit uniqueness via a weaker, but less clean, common belief foundation.

## 4.2 The Common Prior Assumption

We focussed on the case where the common prior assumption holds. Under the common prior assumption, if there is common certainty that players have rank belief  $p$ , i.e., they assign probability  $p$  to having the higher rank, then it must be the case that  $p = \frac{1}{2}$ . On the other hand, without the common prior assumption, common certainty of alternative rank beliefs is possible. Our uniqueness results extend immediately to common certainty of non-uniform rank beliefs, although the nature of global game selection is then different.

Specifically, define  $RB_\varepsilon^p$  for the set of type profiles  $(t_1, t_2)$  where both players have  $\varepsilon$ -rank belief  $p$ :

$$RB_\varepsilon^p = \{(t_1, t_2) \mid p - \varepsilon \leq r_i(t_i) \leq \bar{r}_i(t_i) \leq p + \varepsilon \text{ for each } i\}.$$

Say that action invest is  $p$ -dominant for  $t_i$  if

$$x_i(t_i) \geq p.$$

We write  $D^p$  for the set of type profiles for which invest is  $p$ -dominant for each player, so

$$D^p = \{(t_1, t_2) \mid x_i(t_i) \geq p \text{ for each } i\}.$$

**Proposition 5** *Invest is the uniquely rationalizable action for both players if it is  $p$ -dominant for both players and there is common  $(1 - \varepsilon)$ -belief of  $\varepsilon$ -rank belief  $p + 2\varepsilon$ , i.e., if*

$$(t_1, t_2) \in D^p \cap C^{1-\varepsilon, 1-\varepsilon}(RB_\varepsilon^{p+2\varepsilon}).$$

This observation, suitably generalized to many players, is the driving force behind the results of Izmailkov and Yildiz (2010).

### 4.3 Many Players

We focussed on the case of two players. The extension of the results of this paper to  $N$  players—maintaining the symmetry and separability of payoffs assumptions in this paper—is straightforward, and we describe this extension in this section.

Suppose that each player  $i$  had a payoff type  $x_i$ , the payoff to not investing was 0, and the payoff to investing was  $x_i - 1 + \psi(l)$  where  $l$  is the proportion of other players investing and  $\psi : [0, 1] \rightarrow [0, 1]$  is increasing with  $\psi(0) = 0$  and  $\psi(1) = 1$ . In the special case of two players, these payoffs reduce to those studied in this paper.

We can give a common belief characterization of rationalizability in this setting, although the relevant belief operator is more complicated than in the two player case. For a fixed simple event  $E$ , agent  $i$  and integer  $n$ , we can consider derived events corresponding the set of type profiles  $t_{-i}$  of players other than  $i$  where  $t_j \in E_j$  for exactly  $n$  out of those other players. For each  $n$ , we will be interested in the probability that player  $i$  assigns to such derived events. We will now be interested in when the weighted sum of these probabilities is above some type-dependent level. Thus we define generalized belief operators:

$$B_i^{f_i}(E) = \left\{ t \mid t_i \in E_i \text{ and } \sum_{n=0}^{N-1} \psi\left(\frac{n}{N-1}\right) \pi_i(\{t_{-i} \mid \#\{j \mid t_j \in E_j\} = n\} \mid t_i) \geq f_i(t_i) \right\}.$$

Note that this is a generalization of the type-dependent belief operators that we introduced in the two player case. It is qualitatively more complicated, though, because it depends on the weighted sum of probabilities assigned to a set of events (rather than probabilities of one event). However, with these operators, we can generalize the tight characterization of rationalizability. For a vector of type dependent probability functions  $f = (f_1, \dots, f_N)$ , we can now define  $f$ -belief and common  $f$ -belief operators as before,

$$B_*^f(E) = \bigcap_{i=1}^N B_i^{f_i}(E).$$

$$C^f(E) = \cap_{n=1}^{\infty} [B^f]^n(E).$$

and analogous fixed point characterizations will hold. Now Lemma 2 will continue to hold as stated for these modified operators, as will Proposition 1 giving a sufficient condition for a uniquely rationalizable action in symmetric games. It is also possible to give generalized belief operator characterizations of rationalizable actions in more general games, see Morris and Shin (2007).

While the tight characterization of rationalizability becomes more complicated when we move to many players, the sufficient conditions we give for limit multiplicity and limit uniqueness generalize straightforwardly.

The approximate common certainty sufficient condition for multiplicity in Proposition 2 is sufficient to show rationalizability of all strict equilibrium actions in general games, as originally shown by Monderer and Samet (1989), and thus the fat tails limit multiplicity result of Corollary 6 holds for general games also.

For uniqueness sufficient conditions, a player's rank belief now gives the probability that he assigns to his payoff type being ranked  $k$ th for each  $k$ , and we can define corresponding upper and lower rank beliefs. Thus

$$\begin{aligned}\bar{r}_i(k|t_i) &= \pi_i(\#\{t_j|x_j(t_j) \leq x_i(t_i)\} = k|t_i), \\ \underline{r}_i(k|t_i) &= \pi_i(\#\{t_j|x_j(t_j) < x_i(t_i)\} = k|t_i).\end{aligned}$$

We say that rank belief of a type  $t_i$  is  $\varepsilon$ -uniform if

$$\frac{1}{N} - \varepsilon \leq \underline{r}_i(k|t_i) \leq \bar{r}_i(k|t_i) \leq \frac{1}{N} + \varepsilon.$$

for each  $k = 1, \dots, N$ . We write  $URB_\varepsilon$  for the set of type profiles  $t$  where both players have  $\varepsilon$ -uniform rank beliefs:

$$URB_\varepsilon = \left\{ t \mid \frac{1}{N} - \varepsilon \leq \underline{r}_i(n|t_i) \leq \bar{r}_i(n|t_i) \leq \frac{1}{N} + \varepsilon \text{ for each } i \right\}.$$

We say that rank beliefs are *approximately uniform* if they are  $\varepsilon$ -uniform for some  $\varepsilon \geq 0$ .

Morris and Shin (2003b) noted that the global game selection in this case was the "Laplacian action", corresponding to a uniform belief over the proportion of opponents investing. Our second concept is a strict version of the Laplacian property. We say that action invest is  $\varepsilon$ -Laplacian for  $t_i$  if

$$x_i(t_i) \geq 1 - \frac{1}{N} \sum_{j=0}^{N-1} \psi\left(\frac{j}{N-1}\right) + \varepsilon.$$

We write  $L_\varepsilon$  for the set of type profiles for which invest is  $\varepsilon$ -strictly Laplacian for each player, so

$$L_\varepsilon = \left\{ t \mid x_i(t_i) \geq 1 - \frac{1}{N} \sum_{j=0}^{N-1} \psi\left(\frac{j}{N-1}\right) + \varepsilon \text{ for each } i \right\}.$$

**Proposition 6** *Invest is the uniquely rationalizable action for both players if it is  $\varepsilon$ -Laplacian for both players and there is common  $(1 - \varepsilon)$ -belief of  $\varepsilon$ -uniform rank beliefs for some  $\varepsilon \geq 0$ , i.e., if*

$$t \in L_\varepsilon \cap C^{1-\varepsilon}(URB_\varepsilon).$$

This result easily implies the fat tails limit uniqueness result of Corollary 7, since sufficient conditions for  $\varepsilon$ -uniform rank beliefs in the two player case also imply  $\varepsilon$ -uniform rank beliefs in the many player case.

Two papers that use explicit statements about higher-order beliefs to give sufficient conditions for unique rationalizable actions are Morris and Shin (2012) and Morris (2014). Morris and Shin (2012) consider a particular coordination problem that arises among uninformed investors deciding whether to trade in a market with adverse selection. They say that there is "market confidence" if statements of the form "most players think that..." ( $k$  times) that the losses from informed trade are sufficiently small are true. This corresponds to a many player example of the higher-order belief characterization of Proposition 1. Morris (2014) discusses higher-order beliefs sufficient conditions for coordination when there is synchronization problem. Oyama and Takahashi (2013) use generalized belief operators as tool to prove results about robustness to incomplete information games in the sense of Kajii and Morris (1997).

#### 4.4 Private Values and Separable Payoffs

We simplified the analysis by focussing on a private value global game rather than the more often studied common value case. The alternative "common value" case would correspond to the case where player  $i$ 's payoff is a function of  $y + \tau\eta$ , i.e., the prior mean plus the common shock, but he still observes a signal  $t_i = \tau\eta + \sigma e_i$  which contains an idiosyncratic (but now payoff irrelevant) noise. Much of the literature on global games has focussed on the common value case (e.g., Carlsson and van Damme (1993), Morris and Shin (1998) and Frankel, Morris, and Pauzner (2003)). However, the private value case (Morris and Shin (2005) and Argenziano (2008)) is as tractable and more relevant for many applications.

One complication of extending the analysis to the common value case is that, with regularly varying tails, the underlying payoff functions will no longer satisfy increasing differences, so that higher types of a player have a greater incentive to invest given the other players' action. In particular, in the fat tails case, the distribution of  $\theta = y + \tau\eta$  conditional on  $t_i = \tau\eta + \sigma e_i$  need not be monotone with respect to  $t_i$ .

An additional simplifying assumption we made was that payoffs were additively separable between a component that depended on the opponent's action and a component that depended on an unknown payoff state. This additive separability allows for a tighter description of the connection. Extensions are possible here also, but are messy and involve tedious continuity arguments. See Morris and Shin (2007).

## 4.5 Asymmetric Payoffs

Like much of the applied literature, we focussed on a game which was symmetric across players. However, global game results go through with asymmetric games. One can state analogous higher-order belief properties driving global game results (relating to translation invariance) but they are not as clean. The issue is discussed in Morris and Shin (2007).

## 4.6 Global Games Literature

Let us briefly summarize how the results described in this note relate to the existing global games literature.

The classical exercise in the global games literature initiated by Carlsson and van Damme (1993) is to consider what happens if we fix a prior over payoff relevant states and let the size of the noise in players' conditionally independent signals of the state converge to zero. A key step in such arguments (e.g., in Carlsson and van Damme (1993) and Frankel, Morris, and Pauzner (2003)) is the assumption that the prior distribution is smooth, which ensures that conditional probabilities converge uniformly, or uniformly over a compact interval. In the symmetric case, globally uniform convergence of conditional probabilities gives common certainty of rank beliefs. However, as highlighted by lemma 2, it is enough to have common certainty of rank beliefs on a compact interval including states where players actions are undominated. In this sense, we highlight in this note (using a special case) the properties of higher-order beliefs that drive results in the symmetric case.

Another exercise in the global games literature is to assume that both the prior and private signals are normally distributed and see what happens when both prior distribution and distribution of noise in private signals shrink to zero (Morris and Shin (2001) and Morris and Shin (2003a)). We discussed this case in section 4.1. In this case, one can explicitly compute rank beliefs and see how they evolve as the distributions shrink to zero at different rates. However, this paper has highlighted the fact that these results are special. Focussing on the most interesting case of fat tailed distributions, we have identified distinct properties that give limit uniqueness and limit multiplicity depending on the tail properties of the distributions.

Our results generalize those in the existing literature because we do not exploit monotonicity properties of the type space and thus the game is not supermodular. This is true both with general type spaces, or with one dimensional type spaces with fat-tailed distributions. An order structure on types is used in the existing literature. In the analysis of Carlsson and van Damme (1993), Morris and Shin (2003b) and Frankel, Morris, and Pauzner (2003), there is monotonicity with respect to types in the limit (as noise goes to zero) and continuity arguments are used to provide results in the (not necessarily) monotonic type space away from the limit. In the normal models of Morris and Shin (2001) and Morris and Shin (2003a), monotonicity away from the limit is implied by the normal distribution. In highlighting the connection



between general supermodular games and global games, van Zandt and Vives (2007) impose monotonicity away from the limit in general games. Mathevet (2010) imposes a stochastic dominance property even in the limit to prove global game uniqueness results via a contraction argument. By contrast, results in this paper are proved without any order structure on types.

## A Omitted Proofs

**Proof of Lemma 4.** Take any  $\varepsilon > 0$ . We will find  $\bar{n} < \infty$  such that for all  $n > \bar{n}$  and  $t \in [0, \bar{t}]$ ,<sup>7</sup>  $\mu_n(\cdot|t)$  puts at most probability  $\varepsilon$  on  $(-\infty, -\varepsilon)$  and  $(\varepsilon, \infty)$ , for which  $\mu_\infty(\cdot|t)$  assigns zero probability. Furthermore, when  $t > 2\varepsilon$ ,  $\mu_n(\cdot|t)$  puts at most probabilities  $1 - q(t) + \varepsilon$ ,  $\varepsilon$ , and  $q(t) + \varepsilon$  on sets  $[-\varepsilon, \varepsilon]$ ,  $[\varepsilon, t - \varepsilon]$ , and  $[t - \varepsilon, t + \varepsilon]$ , respectively. Consequently  $\mu_n(\cdot|t)$  puts at least probabilities  $1 - q(t) - 4\varepsilon$  and  $q(t) - 4\varepsilon$  on sets  $[-\varepsilon, \varepsilon]$  and  $[t - \varepsilon, t + \varepsilon]$ , respectively, for which  $\mu_\infty(\cdot|t)$  assigns probabilities  $1 - q(t)$  and  $q(t)$ , respectively.

Now,  $F(\varepsilon/\sigma_n)$  and  $G(\varepsilon/\tau_n)$  converge to 1, while  $F(-\varepsilon/\sigma_n) = 1 - F(\varepsilon/\sigma_n)$  and  $G(-\varepsilon/\tau_n) = 1 - G(\varepsilon/\tau_n)$  converge to 0. Hence, there exists  $\bar{n}_1 < \infty$  such that

$$\varepsilon > \max \left\{ \frac{F(-\varepsilon/\sigma_n)}{F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)}, \frac{G(-\varepsilon/\tau_n)}{G(\varepsilon/\tau_n) - G(-\varepsilon/\tau_n)} \right\} \quad (12)$$

for all  $n > \bar{n}_1$ .

Consider any  $t \geq 0$  and any  $n > \bar{n}_1$ . We start by showing that  $\mu_n(\cdot|t)$  puts at most probability  $\varepsilon$  on  $(-\infty, -\varepsilon)$ :

$$\begin{aligned} \mu_n(\tau_n \eta < -\varepsilon|t) &\leq I_n(t + \varepsilon, \infty) / I_n(t - \varepsilon, t + \varepsilon) \\ &\leq \frac{\tau_n G(-\varepsilon/\tau_n) f((t + \varepsilon)/\sigma_n)}{\tau_n (G(\varepsilon/\tau_n) - G(-\varepsilon/\tau_n)) f((t + \varepsilon)/\sigma_n)} \\ &= \frac{G(-\varepsilon/\tau_n)}{G(\varepsilon/\tau_n) - G(-\varepsilon/\tau_n)} < \varepsilon. \end{aligned}$$

Here, the first inequality is by definition and the monotonicity of  $I_n$ , while the last inequality is by (12). To see the second inequality, note that

$$I_n(t + \varepsilon, \infty) = \int_{-\infty}^{-\varepsilon/\tau_n} f((t - \tau_n \eta)/\sigma_n) g(\eta) \tau_n d\eta \leq \tau_n G(-\varepsilon/\tau_n) f((t + \varepsilon)/\sigma_n),$$

and (similarly)

$$I_n(t - \varepsilon, t + \varepsilon) \geq (G(\varepsilon/\tau_n) - G(-\varepsilon/\tau_n)) \tau_n f\left(\frac{t + \varepsilon}{\sigma_n}\right).$$

Using symmetric arguments, we next conclude that  $\mu_n(\cdot|t)$  puts at most probability  $\varepsilon$  on  $(t + \varepsilon, \infty)$ :

$$\begin{aligned} \mu_n(\tau_n \eta > t + \varepsilon|t) &\leq I_n(-\infty, -\varepsilon) / I_n(-\varepsilon, \varepsilon) \\ &\leq \frac{\sigma_n F(-\varepsilon/\sigma_n) g((t + \varepsilon)/\tau_n)}{\sigma_n (F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)) g((t + \varepsilon)/\tau_n)} < \varepsilon. \end{aligned}$$

When  $t \leq 2\varepsilon$ , this is sufficient because it shows that both  $\mu_n(\cdot|t)$  and  $\mu_\infty(\cdot|t)$  put at least probability  $1 - 2\varepsilon$  on  $[-3\varepsilon, 3\varepsilon]$ .

<sup>7</sup>Since  $f$  and  $g$  are symmetric, it suffices to consider the case  $z \geq 0$ .

Now consider any  $t > 2\varepsilon$ . For this case, we will further show that  $\mu_n(\cdot|t)$  and  $\mu_\infty(\cdot|t)$  assign similar probabilities on sets  $[-\varepsilon, \varepsilon]$ ,  $[\varepsilon, t - \varepsilon]$ , and  $[t - \varepsilon, t + \varepsilon]$ . Firstly, for any  $n > \bar{n}_1$ ,

$$\begin{aligned} \mu_n(\varepsilon < \tau_n \eta < t - \varepsilon|t) &\leq I_n(\varepsilon, t - \varepsilon) / I_n(-\varepsilon, \varepsilon) \\ &\leq \frac{\sigma_n (1 - F(\varepsilon/\sigma_n)) g(\varepsilon/\tau_n)}{\sigma_n (F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)) g((t + \varepsilon)/\tau_n)} \\ &= \frac{F(-\varepsilon/\sigma_n)}{F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)} \frac{g(\varepsilon/\tau_n)}{g((t + \varepsilon)/\tau_n)} \\ &\leq \frac{F(-\varepsilon/\sigma_n)}{F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)} \frac{g(\varepsilon/\tau_n)}{g((\bar{t} + \varepsilon)/\tau_n)}. \end{aligned}$$

Here, the equality is by  $1 - F(\varepsilon/\sigma_n) = F(-\varepsilon/\sigma_n)$ , and the last inequality is by monotonicity of  $g$ . To see the inequality on the second line, note that

$$I_n(\varepsilon, t - \varepsilon) = \int_\varepsilon^{t-\varepsilon} f(z/\sigma_n) g((t - z)/\tau_n) dz \leq g(\varepsilon/\tau_n) \int_\varepsilon^{t-\varepsilon} f(z/\sigma_n) dz;$$

the lower bound for  $I_n(-\varepsilon, \varepsilon)$  has been established in the previous case. Now, since  $g$  has regularly-varying tails,  $g(\varepsilon/\tau_n)/g((\bar{t} + \varepsilon)/\tau_n) \rightarrow (\varepsilon/(\bar{t} + \varepsilon))^{-\beta} < \infty$ . Hence, there exists  $\bar{n}_2 > \bar{n}_1$  such that

$$\frac{F(-\varepsilon/\sigma_n)}{F(\varepsilon/\sigma_n) - F(-\varepsilon/\sigma_n)} \frac{g(\varepsilon/\tau_n)}{g((\bar{t} + \varepsilon)/\tau_n)} < \varepsilon \quad (\forall n > \bar{n}_2).$$

Therefore,

$$\mu_n(\varepsilon < \theta < t - \varepsilon|t) < \varepsilon \quad (\forall n > \bar{n}_2, t \in [2\varepsilon, \bar{t}]).$$

(Note that  $\bar{n}_2$  does not depend on  $t$ .)

Bounding the probabilities of the sets  $[-\varepsilon, \varepsilon]$  and  $[t - \varepsilon, t + \varepsilon]$  requires more subtle arguments. To this end, we construct sequences  $e_n = \sigma_n^{-\gamma}$  and  $\eta_n = \tau_n^{-\gamma}$  for some  $\gamma$  with  $\max\{1/\alpha, 1/\beta\} < \gamma < 1$ . Since  $(\sigma_n, \tau_n) \rightarrow (0, 0)$ , one can take  $e_n > \bar{e}$  and  $\eta_n > \bar{\eta}$  without loss of generality. Moreover,  $\sigma_n e_n \rightarrow 0$ ,  $\tau_n \eta_n \rightarrow 0$ ,  $e_n \rightarrow \infty$ , and  $\eta_n \rightarrow \infty$ , and we consider only  $\sigma_n e_n < \varepsilon$  and  $\tau_n \eta_n < \varepsilon$ . Now,

$$\mu_n(t - \varepsilon < \theta < t + \varepsilon|t) = I_n(-\varepsilon, \varepsilon) / I_n(-\infty, \infty).$$

Towards finding an upper bound, we first obtain the following tighter lower bound for  $I_n(-\infty, \infty)$ :

$$\begin{aligned} I_n(-\infty, \infty) &> I_n(-\sigma_n e_n, \sigma_n e_n) + I_n(t - \tau_n \eta_n, t + \tau_n \eta_n) \\ &\geq (F(e_n) - F(-e_n)) \sigma_n g\left(\frac{t + \sigma_n e_n}{\tau_n}\right) + (G(\eta_n) - G(-\eta_n)) \tau_n f\left(\frac{t + \tau_n \eta_n}{\sigma_n}\right) \equiv D_n(t). \end{aligned}$$

We next find the following upper bounds:

$$\begin{aligned} I_n(-\varepsilon, -\sigma_n e_n) &\leq (F(-e_n) - F(-\varepsilon/\sigma_n)) \sigma_n g\left(\frac{t + \sigma_n e_n}{\tau_n}\right) \leq F(-e_n) \sigma_n g\left(\frac{t + \sigma_n e_n}{\tau_n}\right); \\ I_n(\sigma_n e_n, \varepsilon) &\leq F(-e_n) \sigma_n g\left(\frac{t - \varepsilon}{\tau_n}\right); \\ I_n(-\sigma_n e_n, \sigma_n e_n) &\leq (F(e_n) - F(-e_n)) \sigma_n g\left(\frac{t - \sigma_n e_n}{\tau_n}\right). \end{aligned}$$

Combining with the lower bound above, we further observe that

$$\begin{aligned} I_n(-\varepsilon, -\sigma_n e_n) / D_n(t) &\leq \frac{F(-e_n)}{F(e_n) - F(-e_n)}; \\ I_n(\sigma_n e_n, \varepsilon) / D_n(t) &\leq \frac{F(-e_n)}{F(e_n) - F(-e_n)} \frac{g\left(\frac{t-\varepsilon}{\tau_n}\right)}{g\left(\frac{t+\sigma_n e_n}{\tau_n}\right)}; \\ I_n(-\sigma_n e_n, \sigma_n e_n) / D_n(t) &\leq \frac{(F(e_n) - F(-e_n)) \sigma_n g\left(\frac{t-\sigma_n e_n}{\tau_n}\right)}{(F(e_n) - F(-e_n)) \sigma_n g\left(\frac{t+\sigma_n e_n}{\tau_n}\right) + (G(\eta_n) - G(-\eta_n)) \tau_n f\left(\frac{t+\tau_n \eta_n}{\sigma_n}\right)}. \end{aligned}$$

Now, clearly, the upper bound for  $I_n(-\varepsilon, -\sigma_n e_n) / D_n(t)$  converges to 0. Since  $g\left(\frac{t-\varepsilon}{\tau_n}\right) / g\left(\frac{t+\sigma_n e_n}{\tau_n}\right) \rightarrow \left(\frac{t-\varepsilon}{t}\right)^{-\beta} < 2^\beta$ , the upper bound for  $I_n(\sigma_n e_n, \varepsilon) / D_n(t)$  also goes to 0. Finally, since  $g\left(\frac{t-\sigma_n e_n}{\tau_n}\right) / g\left(\frac{t}{\tau_n}\right)$ ,  $g\left(\frac{t+\sigma_n e_n}{\tau_n}\right) / g\left(\frac{t}{\tau_n}\right)$ , and  $f\left(\frac{t+\tau_n \eta_n}{\sigma_n}\right) / f\left(\frac{t}{\sigma_n}\right)$  all uniformly converge to 1, the upper bound for  $I_n(-\sigma_n e_n, \sigma_n e_n) / D_n(t)$  uniformly converges to  $q(t)$ . Here, all the limits are taken uniformly over  $t \in [2\varepsilon, \bar{t}]$ , and hence there exists  $\bar{n}_3$  such that

$$\mu_n(t - \varepsilon < \tau_n \eta < t + \varepsilon | t) \leq \frac{I_n(-\varepsilon, \varepsilon)}{D_n(t)} < q(t) + \varepsilon \quad (\forall n > \bar{n}_3, t \in [2\varepsilon, \bar{t}]).$$

It is important to observe that  $\bar{n}_3$  does not depend on  $t$ . Similarly, one can find  $\bar{n}_4$  such that

$$\mu_n(-\varepsilon < \tau_n \eta < \varepsilon | t) < 1 - q(t) + \varepsilon \quad (\forall n > \bar{n}_3, t \in [2\varepsilon, \bar{t}]).$$

Setting  $\bar{n} = \max\{\bar{n}_1, \bar{n}_2, \bar{n}_3, \bar{n}_4\}$ , one concludes the proof. ■

**Proof of Lemma 5.** We will consider events  $C \subseteq \mathbb{R}$  corresponding to common shock  $c = \tau \eta$ . For every event  $C$ , define  $r_{n,C} : \mathbb{R} \rightarrow [0, 1]$  by

$$r_{n,C}(t) \equiv \int_C F((t-c)/\sigma_n) d\mu_n(c|t).$$

Note that  $\Pr(x_j < x_i | t_i, \sigma_n, \tau_n, c) = F((t_i - c)/\sigma_n)$ , and hence

$$r_{\sigma_n, \tau_n}(t) = \int F((t-c)/\sigma_n) d\mu_n(c|t) = r_{n,(-\infty, \infty)}(t).$$

Towards computing the limit of  $r_{n,(-\infty, \infty)}$ , we fix  $\varepsilon \in (0, \underline{z})$  and partition  $(-\infty, \infty)$  into intervals

$$\begin{aligned} C_1 &= \mathbb{R} \setminus (t - \varepsilon, t + \varepsilon), \\ C_{2,n} &= (t - \varepsilon, t - \sigma_n e_n) \cup (t + \sigma_n e_n, t + \varepsilon), \\ C_{3,n} &= [t - \sigma_n e_n, t + \sigma_n e_n], \end{aligned}$$

using the notation in the proof of Lemma 4.

We start with  $C_1$ . For each value  $c$  of common shock, define  $F_\infty(c|t) \equiv \lim_n F((t-c)/\sigma_n)$  and observe that it is 1 when  $t > c$  and 0 when  $t < c$ . Since  $F_\infty(c|t)$  is continuous on  $C_1$ ,

$$\lim_{n \rightarrow \infty} r_{n,C_1}(t) = \int_{C_1} F_\infty(c|t) d\mu_\infty(c|t) = F_\infty(0|t) (1 - q(t)).$$

For  $C_{2,n}$ , since  $F(\cdot|t) \in [0, 1]$ , we write

$$0 \leq r_{n,C_{2,n}}(t) \leq \mu_n(C_{2,n}|t)$$

and observe from the proof of Lemma 4 that  $\mu_n(C_{2,n}|t)$  uniformly converges to zero. Finally, we obtain bounds for  $r_{n,C_{3,n}}(t)$ . For the upper bound, we write

$$\begin{aligned} r_{n,C_{3,n}}(t) &\leq \int_{-\sigma_n e_n}^{\sigma_n e_n} F(z/\sigma_n) f(z/\sigma_n) g((t-z)/\tau_n) dz / D_n(t) \\ &\leq g\left(\frac{t - \sigma_n e_n}{\tau_n}\right) \int_{-\sigma_n e_n}^{\sigma_n e_n} F(z/\sigma_n) f(z/\sigma_n) dz / D_n(t) \\ &= \frac{1}{2} (F(e_n) - F(-e_n)) g\left(\frac{t - \sigma_n e_n}{\tau_n}\right) / D_n(t). \end{aligned}$$

Similarly, for the lower bound, we write

$$r_{n,C_{3,n}}(t) \geq \frac{1}{2} (F(e_n) - F(-e_n)) g\left(\frac{t + \sigma_n e_n}{\tau_n}\right) / D_n(t).$$

As in the proof of Lemma 4, both bounds uniformly converge to  $q(t)/2$ . Therefore,

$$r_{\sigma_n, \tau_n}(t) = r_{n,C_1}(t) + r_{n,C_{2,n}}(t) + r_{n,C_{3,n}}(t)$$

uniformly converges to

$$\begin{aligned} r_\infty(t) &= F_\infty(0|t) (1 - q(t)) + q(t) / 2 \\ &= \begin{cases} 1 - q(t) / 2 & \text{if } t > 0 \\ q(t) / 2 & \text{if } t < 0. \end{cases} \end{aligned}$$

■

## References

- ACEMOGLU, D., V. CHERNOZHUKOV, AND M. YILDIZ (2006): “Learning and Disagreement in an Uncertain World,” Discussion paper, M.I.T.
- ANGELETOS, G.-M., AND I. WERNING (2006): “Crises and Prices: Information, Multiplicity and Volatility,” *American Economic Review*, 96, 1720–1736.
- ANGELETOS, M., C. HELLWIG, AND A. PAVAN (2007): “Dynamic Global Games of Regime Change: Learning, Multiplicity and Timing of Attacks,” *Econometrica*, 75, 711–756.
- ARGENZIANO, R. (2008): “Differentiated networks: Equilibrium and Efficiency,” *The RAND Journal Of Economics*, 39, 747–769.
- ATKESON, A. (2001): “Discussion of Morris and Shin,” *NBER Macroeconomics Annual*, 2000, 161–164.
- AUMANN, R. (1976): “Agreeing to Disagree,” *Annals of Statistics*, 4, 1236–1239.
- CARLSSON, H., AND E. VAN DAMME (1993): “Global Games and Equilibrium Selection,” *Econometrica*, 61, 989–1018.
- FRANKEL, D., S. MORRIS, AND A. PAUZNER (2003): “Equilibrium Selection in Global Games with Strategic Complementarities,” *Journal of Economic Theory*, 108, 1–44.

- HEIFETZ, A., AND Z. NEEMAN (2006): “On the Generic (Im)Possibility of Full Surplus Extraction in Mechanism Design,” *Econometrica*, 74, 213–233.
- HELLWIG, C. (2002): “Public Information, Private Information, and the Multiplicity of Equilibria in Coordination Games,” *Journal of Economic Theory*, 107, 191–222.
- HELLWIG, C., A. MUKERJI, AND A. TSYVINSKI (2006): “Self-Fulfilling Currency Crises: The Role of Interest Rates,” *American Economic Review*, 96, 1769–1787.
- IZMALKOV, S., AND M. YILDIZ (2010): “Investor Sentiments,” *American Economic Journal: Microeconomics*, 2, 21–38.
- KAJII, A., AND S. MORRIS (1997): “The Robustness of Equilibria to Incomplete Information,” *Econometrica*, 65, 1283–1309.
- MATHEVET, L. (2010): “A Contraction Principle for Finite Global Games,” *Economic Theory*, 42, 539–563.
- MERTENS, J., AND S. ZAMIR (1985): “Formalization of Bayesian Analysis for Games with Incomplete Information,” *International Journal of Game Theory*, 14, 1–29.
- MONDERER, D., AND D. SAMET (1989): “Approximating Common Knowledge with Common Belief,” *Games and Economic Behavior*, 1, 170–190.
- MORRIS, S. (2014): “Coordination, Timing and Common Knowledge,” *Research in Economics*, 68, 306–314.
- MORRIS, S., AND H. SHIN (1998): “Unique Equilibrium in a Model of Self-Fulfilling Currency Attacks,” *American Economic Review*, 88, 587–597.
- MORRIS, S., AND H. SHIN (2001): “Rethinking Multiple Equilibria in Macroeconomics,” *NBER Macroeconomics Annual*, 2000, 139–161.
- MORRIS, S., AND H. SHIN (2003a): “Coordination Risk and the Price of Debt,” *European Economic Review*, 48, 133–153.
- (2003b): “Global Games: Theory and Applications,” in *Advances in Economics and Econometrics: Proceedings of the Eight World Congress of the Econometric Society*, ed. by M. Dewatripont, L. Hansen, and S. Turnovsky, pp. 56–114. Cambridge University Press, Cambridge.
- (2005): “Heterogeneity and Uniqueness in Interaction Games,” in *The Economy as an Evolving Complex System, III*. Santa Fe Institute.
- MORRIS, S., AND H. SHIN (2012): “Contagious Adverse Selection,” *American Economic Journal: Macroeconomics*, 4, 1–21.
- MORRIS, S., AND H. S. SHIN (2007): “Common Belief Foundations in Global Games,” Discussion paper, <http://www.princeton.edu/~smorris/pdfs/cbf.pdf>.
- OYAMA, D., AND S. TAKAHASHI (2013): “Generalized Belief Operator and the Impact of Small Probability Events on Higher Order Beliefs,” Discussion paper, University of Tokyo and National University of Singapore.

SVENNSON, L. (2006): “The Social Value of Public Information: Morris and Shin (2002) is actually pro transparency, not con.,” *American Economic Review*, 96, 448–452.

VAN ZANDT, T. (2010): “Interim Bayesian Nash Equilibrium on Universal Type Spaces for Supermodular Games,” *Journal of Economic Theory*, 145, 249–263.

VAN ZANDT, T., AND X. VIVES (2007): “Monotone Equilibria in Bayesian Games of Strategic Complementarities,” *Journal of Economic Theory*, 134, 339–360.

WEINSTEIN, J., AND M. YILDIZ (2007): “A Structure Theorem for Rationalizability with Applications to Robust Predictions of Refinements,” *Econometrica*, 75, 365–400.